

## Apple Programs

---

### Major Insects

#### **Aphid eggs, woolly apple aphid**

Natural enemies can be effective on aphids. When 20% of colonies have predators a pesticide application may be delayed or eliminated. Use of pesticides with low toxicity to predators will increase biological control. Product recommendations will be effective on apple aphid, apple grain aphid, and rosy apple aphid, but less so on woolly apple aphid.

#### **Apple rust mite**

Apple rust mite feeds on plant foliage, and in very high numbers, can cause shoots to stop growth prematurely. However, in low to moderate numbers, they are generally regarded as an important and positive part of integrated mite management. Predatory mites can eat apple rust mites when spider mites (a more damaging pest) are scarce, and sustain their populations through the season. Choose pesticides that cause minimal harm to apple rust mite unless 1) populations become very high or 2) large early season populations occur on sensitive cultivars where fruit russetting can occur, such as 'Golden Delicious'. Additional Details

#### **Campylomma**

Campylomma is a sporadic pest of apple, and primarily a beneficial insect (pear psylla predator) on pear. It overwinters in the egg stage on the tree in the bark, and emerges just before and during apple bloom. Large populations require control as soon as they are detected; earlier sprays will do a better job of preventing fruit damage. Pay attention to label restrictions of bloom applications to protect pollinators. Petal-fall sprays will kill nymphs, but prevent little if any fruit damage. Additional Details

#### **Codling moth**

Codling moth is the key pest of pome fruits in the PNW. In general, apples are more susceptible than pears, and fruits with softer flesh are more susceptible to attack. The increasing frequency of a third generation, two have been the norm historically, means that growers must be vigilant throughout the growing season, and be aware of phenology (See WSU Decision Aid system at <https://decisionaid.systems>). Codling moth has a long history of becoming resistant to insecticides, thus rotation of materials with different modes of action (MOA) is highly recommended. Avoid using the same MOA against consecutive generation to minimize this danger. The MOA for each material is listed in the tables. Pheromone mating disruption was registered in 1990, and has since been widely adopted in Washington. Use of mating

disruption is now considered the foundation of an IPM program. Supplementing mating disruption with insecticides may be necessary depending on pressure, and using pheromone traps for monitoring populations will prevent unnecessary applications. Detailed recommendations on pheromone placement and timing of sprays is available. Additional Details

### **Lacanobia fruitworm**

Control sprays should be applied by 1230-1250 degree-days, when only about 10% of the larvae are in their 4th instar. This timing represents the best opportunity to control *Lacanobia subjuncta* with a single insecticide application. During the second generation, 10% of 4th instar is estimated at 3050 degree-days. The best timing for an insecticide application against larvae of the second generation is at 3050 degree-days, but no later than 3150 degree-days.

### **Leafrollers (Pandemis, Obliquebanded)**

Pre-bloom applications of pesticides can be effective and will also conserve natural enemies for leafroller and biological control agents of other pests, such as aphids. If treatments for leafrollers were applied at pink and/or bloom, sampling to determine the density of surviving leafrollers should be completed prior to deciding to apply additional controls at this timing. Most products listed act primarily as stomach poisons versus direct contact to residues, therefore, complete coverage is very important to achieve maximal control. Repeating an application of any product should be based on the leafroller population surviving previous treatments. Use the leafroller models on the WSU Decision Aid System (<https://decisionaid.systems>) for the optimum timing. Additional Details

### **Rosy apple aphid**

Starting at pre-pink monitor 5-10 trees from each block in sensitive varieties. Treatment is justified when more than one cluster per tree is infested. Sprays become progressively less effective as the season advances and leaves curl.

### **San Jose scale**

San Jose scale can be a minor pest if adequately controlled, or escalate into a major problem if not. It primarily infests the trunk and limbs, but scale crawlers will settle on the fruit. Damage to this season's crop may become serious, but ultimately the infestation of wood may cause death of limbs or the entire tree. Oil plus an organophosphate in the delayed dormant spray provide control; if the organophosphate is omitted (oil only), monitor the trees carefully and add one of the listed materials if scale become numerous. Additional Details

### **Shothole borer**

Good sanitation (removing large wood prunings, dead limbs, and woodpiles from the orchard) is the best management tactic. Insecticides are only effective against adults. Beetles begin flying in late April and are active through May. The second generation flight begins in late July or early August. Yellow sticky traps placed on orchard borders will detect adult beetle activity. Spraying the border trees (rows) with high water volumes will protect the remainder of the orchard in many situations where external sources are the primary problem. Additional Details

### **Western tentiform leafminer**

For best results against leafminer, use an adjuvant with abamectin and spinosad. See labels for specific adjuvant recommendations.

### **White apple leafhopper**

Adults fly from late May until frost. Monitor nymphs on the underside of leaves. Egg parasitoid *Anagrus spp.* attacks overwintering and summer eggs. Only control this indirect pest when necessary. Additional Details

### **Woolly apple aphid**

Woolly apple aphid has proven to be one of the most difficult of the aphid pests to control in recent years. The broad-spectrum organophosphates used in previous years are no longer used, and relatively few effective materials remain. This aphid is attacked by many predators (syrphid larvae, lady beetles, lacewings, and earwigs) and a parasitoid, *Aphelinus mali*. These natural enemies may provide control under some circumstances, but biological control may be easily disrupted. Avoid using disruptive pesticides if possible, and if necessary, treat with one of the effective insecticides. See tables. Additional Details

## **Major Diseases**

### **Apple mildew**

Apple and pear powdery mildew is caused by the same fungal species *Podosphaera leucotricha* which overwinters in dormant apple buds, whereas its survival in pear remains unknown. When infected buds break in spring, the fungus produces spores that are rain and wind-spread to infect freshly emerged leaves which are highly susceptible to powdery mildew. Germination and infections are optimal at temperatures between 60F and 78F. Wetness plays a marginal role. The fungus then continues with multi-cycle infections through spring and early summer until the production of new leaves and shoots cease. The fungus is slowed down by the rising temperature (above 82F) as summer progresses. Infection resumes in fall where the pathogen overwinters as ascospores (sexual form) or infected buds. Under high disease pressure and mild summer conditions, the fungus can cause russetting on fruits and therefore reduce quality. While no cultivar is immune, cultivars like Granny Smith, Honeycrisp, Idared and Crimson Crisp are highly susceptible, whereas Golden Delicious is susceptible and Fuji, Gala and Red Delicious are the least susceptible. Mildew management should start before bud break and at green tip stage (to reduce spread of new inoculum) with sulfur-based products and continue every 10 to 14 days until the production of new shoots cease. Fungicides from FRAC groups 3, 7, 11 and 19 are effective and SHOULD be ROTATED throughout the season. In growing, regions where scab is a problem, spray programs used to control the latter will control powdery mildew as well. In organic orchards, sulfur, potassium bicarbonate, and some biopesticides usually provide a good level of control.

### **Apple scab**

Scab, caused by the fungus *Venturia inaequalis*, is a major disease of pome fruit in many growing regions, especially those with high rainfall. Symptoms are gray-brown to blackish lesions on leaves and fruit. Scab risk is low under arid conditions in Central Washington. However, some microclimates in the north of the state can be conducive to scab, and therefore, management is recommended. Where scab is a problem, the fungus overwinter in fallen leaves making leaf removal/incorporation critical to reduce inoculum for the following season. Scab is effectively controlled by the same fungicides sprayed for apple powdery mildew including fungicides from FRAC groups 3, 7 and 11.

### **Botrytis-Gray Mold**

Gray mold, caused by *Botrytis*, is the second most important apple fruit disease and can be the most important disease affecting pear as shown in recent statewide and regional surveys in the Pacific Northwest. Flowers of both crops are susceptible to *Botrytis* infections which persist throughout the growing season until harvest. *Botrytis* infections remain dormant until storage where the fungus causes Gray Mold with symptoms becoming visible after a few months in storage. Afterward, the fungus can spread to healthy fruit. Temperatures between 64F and 78F are optimal for infections. Because infections occur EXCLUSIVELY in the orchard, it is important to start management as early as possible. Delayed management will fail to control infections that started weeks or months before harvest. The fungus is ubiquitous and overwinters on mummified fruit left on trees and fallen leaves. Good sanitation practices will reduce inoculum loads but because of the explosive nature of this disease, fungicide applications are necessary to achieve good control. At bloom time and during spring, fungicides from FRAC groups 7, 9 and 11, used to control apple powdery mildew or scab, will be effective against *Botrytis* if resistance is absent. Fungicides from FRAC 3 have a limited efficacy against *Botrytis* infections. As fruit mature, they become more susceptible to *Botrytis*. Late season management is especially important for cultivars picked after mid-September in WA when wet, disease conducive weather is more likely. Preharvest applications and ROTATIONS of fungicides from the FRAC groups 1, 7, 11 and 19 control *Botrytis*. Tank-mixture of single-site fungicide with Ziram or captan will increase efficacy and delay the selection for resistant populations. IMPORTANT: *Botrytis cinerea* is the MOST RISKY fungus for fungicide resistance development as the fungus can develop resistance to multiple fungicides simultaneously. Remember this aspect when spraying for other diseases such as powdery mildew, as the same fungicides sprayed early in the season can select for resistant *Botrytis* populations which will persist throughout the season and to the storage rooms resulting in limited efficacy of eventual postharvest treatment.

### **Bull's eye rot**

Bull's eye rot is a major disease of apple and pear. The disease can be caused by four different fungal species from the genus *Neofabraea*. The main species causing Bull's eye rot of apple in eastern Washington is *N. perennans*, whereas *N. malicorticis* was reported to be predominant in western Washington. It infects fruit and causes cankers on trees where it overwinters until conditions become favorable in the following spring when it causes new infections. Fruit are infected exclusively in orchards but bull's eye rot symptoms are only seen after several months in cold storage. Therefore, preharvest management is key to reducing decay rates in the packinghouse. Prune cankered branches to reduce the inoculum load and use fungicide applications prior to harvest to control. Ziram applied within two weeks before harvest is recommended for control of Bull's eye rot in the Pacific Northwest. Topsin-M is ONLY recommended under wet conditions and for cultivars, such as Golden Delicious, Pinata, Fuji and Granny Smith, more susceptible to bull's eye rot. Frequent sprays may increase risks of resistance development to Mertect, a fungicide from the same group as Topsin-M (1) used after harvest. Tank-mixtures of Ziram with other single-site fungicides are recommended to increase efficacy and reduced risks of fungicide resistance development.

### **Fire blight**

There is a risk of fire blight infection any time there are flowers on the tree, the weather is warm, and wetting occurs. **Early bloom.** Apply biologicals (Blossom Protect) during early bloom. If fire blight was in the orchard last year apply two applications of the biological. Reapply biological a second time if lime sulfur was applied (Lime sulfur is antimicrobial and kills the biological). **Full bloom to petal fall.** Watch the model. If an infection event occurs, apply an antibiotic as soon as possible, but within

24 hours of the infection (usually wetting of flowers). Repeated antibiotic sprays may be necessary during extended high or extreme risk periods. Best results are obtained when applied within 24-hour window before flower wetting during a high infection risk period. Beneficial only for flower infection prevention. Product used must contact the interior of the flowers in sufficient water and approved wetting agent to completely wet the interior. One pound of any 17% oxytetracycline product per 100 gallons gives a 200 ppm solution. Kasugamycin is another effective antibiotic. Applications of less than 100 gal/A can be effective on small trees if flower interiors are well covered, but do not drop the ppm below 200 (oxytetracycline). Application by ground equipment on each row is highly recommended. Application of antibiotics by aircraft is not recommended. Many fire blight bacteria in the Pacific Northwest are resistant to streptomycin, another registered antibiotic. **Organic. Prebloom: Fixed copper sanitation if fire blight was in the orchard last year. Apples Easy to Thin: Blossom Protect/ Buffer Protect early. a Lime sulfur (+ oil). Second Blossom Protect/ Buffer Protect. Depending on the model and cultivar russet risk soluble copper (Previsto 3 qt, Cueva 4 qt, or Cueva 3 qt + Serenade Opti). Petal fall + 1-2 weeks Serenade Opti (most fruit safe) or 2% lime sulfur (red apples). Apples Hard to Thin/Long Bloom Period: Lime sulfur (+ oil). Blossom Protect + Buffer Protect. a Lime sulfur (+ oil). Second Blossom Protect + Buffer Protect. Depending on the model and cultivar russet risk soluble copper (Previsto 3 qt, Cueva 4 qt, or Cueva 3 qt + Serenade Opti). Petal fall + 1-2 weeks Serenade Opti or 2% lime sulfur (red apples). Apples Hard to thin varieties/ short bloom period: Lime sulfur (+ oil) 2-3 applications. Depending on the model and cultivar russet risk soluble copper (Previsto 3 qt, Cueva 4 qt, or Cueva 3 qt + Serenade Opti).g Petal fall + 1-2 weeks Serenade Opti (most fruit safe) or 2% lime sulfur (red apples). Pears Easy to Mark Varieties: 2 applications of Blossom Protect + Buffer Protect during early bloom to petal fall (70-80% bloom if single treatment).a Follow with Serenade Opti at petal fall to reduce russet risk from Blossom Protect yeast. Pears Marking Tolerant Varieties: 2 applications of Blossom Protect + Buffer Protect during early bloom to petal fall (70-80% bloom if single treatment). Follow with soluble copper (Cueva 4 qrt, Previsto 3 qrt, or Cueva 3 qrt + Serenade Opti) if the model indicates risk (warm/wet).**

#### **Pathogens of Storage Rots**

Several other fungal fruit infections initiated in the orchard can cause storage rots. **Alternaria rot: A dark-brown to black infection caused by *Alternaria alternata* (and other spp.) is ubiquitous in most orchards. Infections, usually sporadic, may become frequent when sanitation is not observed or when wet conditions occur for an extended period. The fungus infects flowers at bloom, but can also infect fruit through the calyx end or wounds. Floral infections can result in moldy-core disease later in storage.**

**Sphaeropsis rot: A sporadic emerging disease caused by the fungus *Sphaeropsis pyriputrescens* infects fruits in the orchard and develops stem and calyx end rots in storage. The fungus overwinters on cankers and twigs. Prune diseased branches to help reduce inoculum. Pruning symptomatic crab apples is particularly important. Although this disease can be sporadic, it is still quarantined in many export countries and its identification in entry ports will result in fruit lot rejection.**

**Yellow-Lambertella rot: This disease was recently reported in the Pacific Northwest and, therefore, is considered as quarantine pathogen. Infections are caused by the fungus *L. corni-maritima*, which has been isolated from mummies of other fruit crops in the past but its disease cycle in apple is still unknown. The disease develops yellow mycelium that cover the fruit, but symptoms are only observed after several months of storage. Recent studies have shown that fungicides from FRAC group 1 are not effective against *Lambertella*, whereas fungicides from FRAC groups 7 and 11 have only moderate efficacy. Until**

*further research has shown which other preharvest fungicides are effective, it is recommended to apply a fungicide from FRAC group 9 or 12 postharvest, as these were found to be the most effective. The fungus requires a wound on the cuticle to cause an infection, therefore, reducing damages and punctures at harvest will reduce infection risks. The possibility of infections occurring through the calyx- or stem-ends is still unknown.*

#### ***Speck rot***

*Speck rot is an emerging disease in the Pacific Northwest and has been recently reported in Europe and Chile. It is caused by the fungus *Phacidiopycnis washingtonensis* that infects fruit exclusively in the orchard. Although its epidemiology is still not fully understood, the primary source of inoculum in the PNW is thought to be Manchurian crab apple pollinizers which are susceptible to the disease. Interestingly, the disease has been reported in other regions where the Manchurian crab apple species are not used. Mummies and dead shoots left on trees from the previous season are the main source of inoculum to infect commercial apple fruit. Therefore, pruning and mummy removal will significantly reduce the inoculum load. Fungicides from FRAC groups 1 and 3 are effective when reduced sensitivity or resistance is absent in the orchard. Both active ingredients of Pristine are not effective against Speck rot. Because the exact time of infections is still unknown, sprays that start a month prior to harvest are recommended.*

## ***Major Fruit Protectants***

#### ***Apple Sunburn***

*Sunburn damage costs apple growers tens of millions of dollars annually and is often the primary cause of fruit cullage for apples grown in the Pacific Northwest. Growers often lose more than 10% of their apples to sunburn unless they have used some means of protecting their fruit from sunburn damage. There are four types of apple sunburn: (i) Sunburn Necrosis, (ii) Sunburn Browning, (iii) Photo-Oxidative Sunburn, and (iv) Storage Sunburn. Apple fruit are susceptible to sunburn because they have a much higher thermal mass (the ability of a material to absorb and store heat energy) than leaves and are not able to dissipate this heat as effectively as leaves. It is important to remember that fruit temperature can be considerably higher (20 - 30°F) than the ambient air temperature.*

# Spray Schedule

## Dormant

Insect	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
San Jose scale	petroleum oil- dormant petroleum oil- dormant	1.5 % v/v	12 h	none listed		4	Organic

Efficacy numbers denote the relative efficacy of a pesticide against a given pest on a 1 to 4 scale with 1 being low and 4 high efficacy. This information is based primarily on research conducted with WSU researchers in Washington.

## Delayed dormant

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
Apple mildew	benzovindiflupyr Aprovia	7 fl oz	12 h	30 d	7	3	Do not follow Aprovia with fungicides from FRAC group 7 such as Fontelis, Luna and Pristine to reduce fungicide resistance development.
	trifloxystrobin Flint Extra	2.9 fl oz	12 h	14 d	11	3	
	penthiopyrad Fontelis	14-20 fl oz	12 h	28 d	7	3	Fontelis is a FRAC 7 fungicide.
	cyprodinil + difenoconazole Inspire Super	12 fl oz	12 h	14 d	9,3	3	Inspire Super contains cyprodinil, an active ingredient similar to the one in Vanguard. Do not rotate this product or use in the same season to avoid fungicide resistance development. Do apply after or before Rally 40WSP or other FRAC group 3 fungicides to minimize fungicide resistance in powdery mildew and other non-target pathogens.
	fluopyram + trifloxystrobin Luna Sensation	5-5.8 fl oz	12 h	14 d	7, 11	4	
	triflumizole Procure 480SC	8-16 fl oz	12 h	14 d	3	4	

<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<b>myclobutanil</b> <b>Rally 40WSP</b>	<b>5 oz</b>	<b>24 h</b>	<b>14 d</b>	<b>3</b>	<b>4</b>	<b>Rally is a FRAC 3 fungicide and should not be rotated or used with other FRAC 3 fungicides. Place Rally into solution before adding oil. Apply no sooner than half-inch green.</b>
	<b>lime sulfur/calcium polysulfide</b> <b>Rex Lime Sulfur</b>	<b>See label</b>	<b>48 h</b>	<b>0 d</b>		<b>NR</b>	<b>This material is toxic to pest and predatory mites; destroying apple rust mites, the alternate prey of predatory mites, may predispose the orchard to later spider mite problems.</b> <span style="border: 1px solid black; padding: 0 2px;">Organic</span>
	<b>flutriafol</b> <b>Topguard</b>	<b>8-12 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>3</b>	<b>4</b>	<b>Topguard is a FRAC group 3 fungicide and should not be rotated or used with other FRAC 3 fungicides. Topguard may cause a mild phytotoxicity on Braeburn apple leaves.</b>
	<b>triflumazole</b> <b>Trionic 4SC</b>	<b>16 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>3</b>	<b>4</b>	<b>Trionic is a FRAC 3 fungicide an should not be rotated or used with other FRAC 3 fungicides.</b>
	<b>Notes: The efficacy level will depend on the absence of resistant populations from the orchard. To limit the potential for development of fungicide resistance, do not make more than four applications of FRAC group 11 fungicides per season. Do not make more than two sequential applications of FRAC group 11 fungicides. This limitation is inclusive of all FRAC group 11 fungicides labeled for use on pome fruits.</b>						
	<b>Notes: To limit the potential for development of fungicide resistance, rotate between FRAC groups. Do not make sequential applications of fungicides in the same FRAC group. Do not make more than 2 applications of fungicides in the same FRAC group per season.</b>						
<b>Apple scab</b>	<b>benzovindiflupyr</b> <b>Aprovia</b>	<b>7 fl oz</b>	<b>12 h</b>	<b>30 d</b>	<b>7</b>	<b>2</b>	
	<b>captan</b> <b>Captan 50WP</b>	<b>6 lb</b>	<b>24 h</b>	<b>0 d</b>	<b>M4</b>	<b>NR</b>	<b>Do not use captan on pink through blossom stages. When possible, tank-mix captan with other single-site fungicides to help reduce fungicide resistance development.</b>
	<b>mancozeb</b> <b>Dithane M45 80W</b>	<b>6 lb</b>	<b>24 h</b>	<b>77 d</b>	<b>M3</b>	<b>NR</b>	<b>Do not apply after bloom. See label for restrictions.</b>
	<b>trifloxystrobin</b> <b>Flint Extra</b>	<b>2.5-2.9 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>11</b>	<b>3</b>	
	<b>penthiopyrad</b> <b>Fontelis</b>	<b>14-20 fl oz</b>	<b>12 h</b>	<b>28 d</b>	<b>7</b>	<b>NR</b>	



<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<i>cyprodinil + difenoconazole Inspire Super</i>	12 fl oz	12 h	14 d	9,3	3	<i>Inspire Super contains cyprodinil, an active ingredient similar to the one in Vanguard. Do not rotate this products or use in the same season to avoid fungicide resistance development. Do apply after or before Rally 40WSP or other FRAC group 3 fungicides to minimize fungicide resistance in powdery mildew and other non-target pathogens.</i>
	<i>fluopyram + trifloxystrobin Luna Sensation</i>	4-5.8 fl oz	12 h	14 d	7, 11	NR	<i>Do not use before or after Fontelis to minimize fungicide resistance development in FRAC group 7. Luna Sensation contains an active ingredient similar to the one in Flint or Sovran. Do not follow-up Luna with either product.</i>
	<i>triflumizole Procure 480SC</i>	8-16 fl oz	12 h	14 d	3	NR	
	<i>myclobutanil Rally 40WSP</i>	5 oz	24 h	14 d	3	NR	<i>Place Rally into solution before adding oil. Apply no sooner than half-inch green.</i>
	<i>lime sulfur/calcium polysulfide Rex Lime Sulfur</i>	See label	48 h	0 d		NR	<i>This material is toxic to pest and predatory mites; destroying apple rust mites, the alternate prey of predatory mites, may predispose the orchard to later spider mite problems. <span style="border: 1px solid black; padding: 0 2px;">Organic</span></i>
	<i>flutriafol Topguard</i>	13 fl oz	12 h	14 d	3	NR	<i>Topguard may cause a mild phytotoxicity on Braeburn apple leaves.</i>
	<i>ziram Ziram 76DF</i>	See label	48 h	14 d	M3	NR	
<i>Notes: To limit the potential for development of fungicide resistance, do not make more than four applications of strobilurin fungicides per season. Do not make more than two sequential applications of strobilurin fungicides. This limitation is inclusive of all strobilurin fungicides (Mode 11) labeled for use on pome fruits.</i>							
<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
<i>Aphid eggs, woolly apple aphid</i>	<i>petroleum oil- dormant petroleum oil- dormant</i>	1.5 % v/v	12 h	none listed		4	<i>Apply chlorpyrifos at half-inch green. If using with oil, liquid formulations are preferred. Do not make more than one application of chlorpyrifos per year. <span style="border: 1px solid black; padding: 0 2px;">Organic</span></i>
<i>Cutworms</i>	<i>indoxacarb Avaunt</i>	See label	12 h	14 d	22	4	
	<i>methoxyfenozide Intrepid 2F</i>	See label	4 h	14 d	18A	NR	

<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
<b>European red mite</b>	<b>petroleum oil- dormant petroleum oil- dormant</b>	<b>1.5 % v/v</b>	<b>12 h</b>	<b>none listed</b>		<b>3-4</b>	<b>Oil is indispensable for an integrated mite control program.</b> <span style="border: 1px solid black; padding: 2px;">Organic</span>
<b>Grape mealybug</b>	<b>diazinon + petroleum oil- dormant Diazinon 50W + petroleum oil- dormant</b>	<b>4 lb 1-1.5 % v/v</b>	<b>4 d</b>	<b>21 d</b>	<b>1B</b>	<b>3</b>	<b>Oil is indispensable for an integrated mite control program.</b>
<b>Lygus and stink bugs</b>	<b>Notes: Do not make more than one application of chlorpyrifos per year.</b>						
<b>San Jose scale</b>	<b>pyriproxyfen + petroleum oil- dormant Esteem 35WP + petroleum oil- dormant</b>	<b>4-5 oz 1-1.5 % v/v</b>	<b>12 h</b>	<b>45 d</b>	<b>7C</b>	<b>1</b>	
	<b>petroleum oil- dormant petroleum oil- dormant</b>	<b>1-1.5 % v/v</b>	<b>12 h</b>	<b>none listed</b>		<b>4</b>	<span style="border: 1px solid black; padding: 2px;">Organic</span>

Efficacy numbers denote the relative efficacy of a pesticide against a given pest on a 1 to 4 scale with 1 being low and 4 high efficacy. This information is based primarily on research conducted with WSU researchers in Washington.

## Prepink

<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
<b>Apple mildew</b>	<b>trifloxystrobin Flint Extra</b>	<b>2.9 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>11</b>	<b>3</b>	
	<b>penthiopyrad Fontelis</b>	<b>14-20 fl oz</b>	<b>12 h</b>	<b>28 d</b>	<b>7</b>	<b>3</b>	<b>Fontelis is a FRAC 7 fungicide.</b>
	<b>cyprodinil + difenoconazole Inspire Super</b>	<b>12 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>9,3</b>	<b>3</b>	<b>Inspire Super contains cyprodinil, an active ingredient similar to the one in Vanguard. Do not rotate this product or use in the same season to avoid fungicide resistance development. Do apply after or before Rally 40WSP or other FRAC group 3 fungicides to minimize fungicide resistance in powdery mildew and other non-target pathogens.</b>
	<b>fluopyram + trifloxystrobin Luna Sensation</b>	<b>5-5.8 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>7, 11</b>	<b>4</b>	
	<b>triflumizole Procure 480SC</b>	<b>8-16 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>3</b>	<b>4</b>	

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
	<i>myclobutanil</i> Rally 40WSP	5 oz	24 h	14 d	3	4	Rally is a FRAC 3 fungicide and should not be rotated or used with other FRAC 3 fungicides. Place Rally into solution before adding oil. Apply no sooner than half-inch green.
	<i>lime sulfur/calcium polysulfide</i> Rex Lime Sulfur	See Label	48 h	0 d		NR	This material is toxic to pest and predatory mites; destroying apple rust mites, the alternate prey of predatory mites, may predispose the orchard to later spider mite problems. <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<i>Bacillus pumilus strain QST 2808</i> Sonata	2-4 qt	4 h	0 d		NR	<span style="border: 1px solid black; padding: 2px;">Organic</span>
	<i>flutriafol</i> Topguard	8-12 fl oz	12 h	14 d	3	4	Topguard is a FRAC group 3 fungicide and should not be rotated or used with other FRAC 3 fungicides. Topguard may cause a mild phytotoxicity on Braeburn apple leaves.
<b>Notes: To limit the potential for development of fungicide resistance, rotate between FRAC groups. Do not make sequential applications of fungicides in the same FRAC group. Do not make more than 2 applications of fungicides in the same FRAC group per season.</b>							
<b>Notes: Apply one Group 3 Fungicide at the same growth stage.</b>							
<b>Notes: The efficacy level will depend on the absence of resistant populations from the orchard. To limit the potential for development of fungicide resistance, do not make more than four applications of FRAC group 11 fungicides per season. Do not make more than two sequential applications of FRAC group 11 fungicides. This limitation is inclusive of all FRAC group 11 fungicides labeled for use on pome fruits.</b>							
<b>Apple scab</b>	<i>captan</i> Captan 50WP	6 lb	24 h	0 d	M4	NR	Do not use captan on pink through blossom stages. When possible, tank-mix captan with other single-site fungicides to help reduce fungicide resistance development.
	<i>mancozeb</i> Dithane M45 80W	6 lb	24 h	77 d	M3	NR	Do not apply after bloom. See label for restrictions.
	<i>penthiopyrad</i> Fontelis	14-20 fl oz	12 h	28 d	7	NR	Fontelis and Luna Sensation are from the same chemical group (7). Apply one of them ONLY at the same growth stage.
	<i>cyprodinil + difenoconazole</i> Inspire Super	12 fl oz	12 h	14 d	9,3	NR	Inspire Super and Vanguard 75WG are from the same chemical group. Use one or the other of these products. These two products should not be used in rotation.
	<i>fluopyram + trifloxystrobin</i> Luna Sensation	4-5.8 fl oz	12 h	14 d	7, 11	NR	Luna Sensation and Fontelis are from the same chemical group (7). Apply one of them ONLY at the same growth stage.

<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<i>triflumizole</i> <b>Procure 480SC</b>	8-16 fl oz	12 h	14 d	3	NR	Apply only one Group 3 Fungicide at the same growth stage.
	<i>myclobutanil</i> <b>Rally 40WSP</b>	5 oz	24 h	14 d	3	NR	Apply only one Group 3 Fungicide at the same growth stage.
	<i>lime sulfur/calcium polysulfide</i> <b>Rex Lime Sulfur</b>	See Label	48 h	0 d		NR	This material is toxic to pest and predatory mites; destroying apple rust mites, the alternate prey of predatory mites, may predispose the orchard to later spider mite problems. <span style="border: 1px solid black; padding: 0 2px;">Organic</span>
	<i>flutriafol</i> <b>Topguard</b>	13 fl oz	12 h	14 d	3	NR	Topguard may cause a mild phytotoxicity on Braeburn apple leaves.
	<i>ziram</i> <b>Ziram 76DF</b>	See label	48 h	14 d	M3	NR	
<p><b>Notes: To limit the potential for development of fungicide resistance, do not make more than four applications of strobilurin fungicides per season. Do not make more than two sequential applications of strobilurin fungicides. This limitation is inclusive of all strobilurin fungicides (Mode 11) labeled for use on pome fruits.</b></p>							
<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
<b>Cutworms</b>	<i>indoxacarb</i> <b>Avaunt</b>	See label	12 h	14 d	22	4	
	<i>methoxyfenozide</i> <b>Intrepid 2F</b>	See label	4 h	14 d	18A	NR	
<b>Grape mealybug</b>	<i>acetamiprid</i> <b>Assail 70WP</b>	3.4 oz	12 h	7 d	4A	1	
	<i>buprofezin</i> <b>Centaur WDG</b>	34.5 oz	12 h	14 d	16	NR	
<b>Leafrollers (Pandemis)</b>	<i>Bacillus thuringiensis subsp. kurstaki</i> <b>DiPel DF</b>	1-2 lb	4 h	0 d	11B2	3	While too early for Obliquebanded leafrollers, this timing is appropriate for Pandemis. Bts are stomach poisons so complete coverage is very important for control. Two or three applications are usually required. Apply when forecasts predict a warm weather pattern, above 65°F, for 3 or more days. <span style="border: 1px solid black; padding: 0 2px;">Organic</span>
<b>Lygus and stink bugs</b>	<i>flonicamid</i> <b>Beleaf 50SG</b>	2.8 oz	12 h	21 d	29	NR	Needs further study. 50% control of adults and 65% control of young nymphs in one WA study in alfalfa [Walsh 2018].
	<i>diazinon</i> <b>Diazinon 50W</b>	4 lb	4 d	21 d	1B	NR	

<i>Insect</i>	<i>Chemical</i>	<i>Rate per Acre</i>	<i>REI</i>	<i>PHI</i>	<i>MOA</i>	<i>Eff.</i>	<i>Notes</i>
	<b><i>sulfoxaflor Transform</i></b>	<b><i>2.25 oz</i></b>	<b><i>24 h</i></b>	<b><i>7 d</i></b>	<b><i>4C</i></b>	<b><i>NR</i></b>	<b><i>Needs further testing. 68% control of adults and 71% control of young nymphs in one WA study in alfalfa [Walsh 2018].</i></b>

*Efficacy numbers denote the relative efficacy of a pesticide against a given pest on a 1 to 4 scale with 1 being low and 4 high efficacy. This information is based primarily on research conducted with WSU researchers in Washington.*

---

Pink

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
Apple mildew	benzovindiflupyr Aprovia	5.5-7 fl oz	12 h	30 d	7	3	Do not follow Aprovia with fungicides from FRAC group 7 such as Fontelis, Luna and Pristine to reduce fungicide resistance development.
	trifloxystrobin Flint Extra	2.9 fl oz	12 h	14 d	11	3	
	penthiopyrad Fontelis	14-20 fl oz	12 h	28 d	7	3	Fontelis is a FRAC 7 fungicide.
	cyprodinil + difenoconazole Inspire Super	12 fl oz	12 h	14 d	9,3	3	Inspire Super contains cyprodinil, an active ingredient similar to the one in Vanguard. Do not rotate this product or use in the same season to avoid fungicide resistance development. Do apply after or before Rally 40WSP or other FRAC group 3 fungicides to minimize fungicide resistance in powdery mildew and other non-target pathogens.
	triflumizole Procure 480SC	8-16 fl oz	12 h	14 d	3	4	
	myclobutanil Rally 40WSP	5 oz	24 h	14 d	3	4	Rally is a FRAC 3 fungicide and should not be rotated or used with other FRAC 3 fungicides. Place Rally into solution before adding oil. Apply no sooner than half-inch green.
	Bacillus pumilus strain QST 2808 Sonata	2-4 qt	4 h	0 d		NR	<span style="border: 1px solid black; padding: 2px;">Organic</span>
	lime sulfur/calcium polysulfide Sulforix	See label	48 h	none listed		NR	Toxic to rust mites, the primary alternate prey of predatory mites. Also toxic to spider mites and predatory mites.
	flutriafol Topguard	8-12 fl oz	12 h	14 d	3	NR	Topguard may cause a mild phytotoxicity on Braeburn apple leaves.
Notes: To limit the potential for development of fungicide resistance, rotate between FRAC groups. Do not make sequential applications of fungicides in the same FRAC group. Do not make more than 2 applications of fungicides in the same FRAC group per season.							
Notes: To limit the potential for development of fungicide resistance, do not make more than four applications of strobilurin fungicides per season. Do not make more than two sequential applications of strobilurin fungicides. This limitation is inclusive of all strobilurin fungicides (Mode 11) labeled for use on pome fruits.							

<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
<b>Apple scab</b>	<b>benzovindiflupyr</b> <b>Aprovia</b>	5.5-7 fl oz	12 h	30 d	7	3	<i>Do not follow Aprovia with fungicides from FRAC group 7 such as Fontelis, Luna and Pristine to reduce fungicide resistance development.</i>
	<b>mancozeb</b> <b>Dithane M45 80W</b>	6 lb	24 h	77 d	M3	NR	<i>Do not apply after bloom. See label for restrictions.</i>
	<b>trifloxystrobin</b> <b>Flint Extra</b>	2.5-2.9 fl oz	12 h	14 d	11	3	
	<b>penthiopyrad</b> <b>Fontelis</b>	14-20 fl oz	12 h	28 d	7	NR	<i>Fontelis and Aprovia are from the same chemical group (7). Apply one of them ONLY at the same growth stage.</i>
	<b>cyprodinil + difenoconazole</b> <b>Inspire Super</b>	12 fl oz	12 h	14 d	9,3	3	<i>Inspire Super contains cyprodinil, an active ingredient similar to the one in Vanguard. Do not rotate this products or use in the same season to avoid fungicide resistance development. Do apply after or before Rally 40WSP or other FRAC group 3 fungicides to minimize fungicide resistance in powdery mildew and other non-target pathogens.</i>
	<b>metiram</b> <b>Polyram 80DF</b>	See label	24 h	77 d	M3	NR	
	<b>triflumizole</b> <b>Procure 480SC</b>	8-16 fl oz	12 h	14 d	3	NR	<i>Apply only one Group 3 Fungicide at the same growth stage.</i>
	<b>myclobutanil</b> <b>Rally 40WSP</b>	5 oz	24 h	14 d	3	NR	<i>Apply only one Group 3 Fungicide at the same growth stage. See note for Inspire Super.</i>
	<b>lime sulfur/calcium polysulfide</b> <b>Sulforix</b>	See label	48 h	none listed		NR	<i>Toxic to rust mites, the primary alternate prey of predatory mites. Also toxic to spider mites and predatory mites.</i>
	<b>flutriafol</b> <b>Topguard</b>	13 fl oz	12 h	14 d	3	NR	<i>Topguard may cause a mild phytotoxicity on Braeburn apple leaves.</i>
	<b>ziram</b> <b>Ziram 76DF</b>	See label	48 h	14 d	M3	NR	<i>Besides having some activity against scab, Ziram may reduce bull's eye rot inoculum persisting on cankers from the previous season.</i>
<b>Notes: To limit the potential for development of fungicide resistance, do not make more than four applications of strobilurin fungicides per season. Do not make more than two sequential applications of strobilurin fungicides. This limitation is inclusive of all strobilurin fungicides (Mode 11) labeled for use on pome fruits.</b>							
<b>Fire blight</b>	<b>Aureobasidium pullulans strains DSM 14940 &amp; 14941</b> <b>Blossom Protect</b>	1.25 lb	4 h	none listed		4	<i>Apply with Buffer Protect. 50% and 80% bloom. Yeasts need 1-2 days before an infection to colonize the flower before bacteria invade to be effective. <span style="border: 1px solid black; padding: 0 2px;">Organic</span></i>

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
	lime sulfur/calcium polysulfide lime sulfur/calcium polysulfide	2 % v/v	48 h	none listed		NR	Early bloom applications plus oil are antimicrobial. 20 and 70% bloom timings. Reapply biologicals after lime sulfur if used.
Notes: Early bloom. Apply biologicals (Blossom Protect) during early bloom. If fire blight was in the orchard last year apply two applications of the biological. Reapply biological a second time if lime sulfur was applied (Lime sulfur is antimicrobial and kills biologicals).							
Insect	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
Apple rust mite	spirodiclofen Envidor 2SC	18 fl oz	12 h	7 d	23	NR	
	fenbutatin oxide Vendex 50WP	1-1.5 lb	48 h	14 d	12B	3-4	
Campylocoma	acetamiprid Assail 70WP	1.7-3.4 oz	12 h	7 d	4A	NR	Use higher rates for high population numbers.
Codling moth	CM pheromone dispensers Isomate-C Plus	See label	none listed	none listed		NR	Install dispensers before first flight (prior to bloom) using the full label rate in the top 2 feet of the canopy. When using aerosol emitters borders should be treated with hand-applied dispensers. <span style="border: 1px solid black; padding: 2px;">Organic</span>
Leafrollers (Pandemis, Obliquebanded)	chlorantraniliprole Altacor	4.5 oz	4 h	5 d	28	4	
	Bacillus thuringiensis subsp. kurstaki DiPel DF	See label	4 h	0 d	11B2	3	Bts are stomach poisons so complete coverage is very important for control. Two or three applications are usually required. Apply when forecasts predict a warm weather pattern, above 65°F, for 3 or more days. <span style="border: 1px solid black; padding: 2px;">Organic</span>
	methoxyfenozide Intrepid 2F	16 fl oz	4 h	14 d	18A	3	
Notes: Target leafroller larvae in the spring at 300-560 DD (Obliquebanded) and 300-460 DD (Pandemis). Target 1st summer generation larvae at 955-1300 DD (Obliquebanded) and 1760-2255 DD (Pandemis).							
Rosy apple aphid	acetamiprid Assail 70WP	1.7 oz	12 h	7 d	4A	3-4	Use higher rates for high population numbers.

Efficacy numbers denote the relative efficacy of a pesticide against a given pest on a 1 to 4 scale with 1 being low and 4 high efficacy. This information is based primarily on research conducted with WSU researchers in Washington.

## Bloom



<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
<b>Apple mildew</b>	<b>polyoxin D zinc salt Ph-D</b>	<b>6.2 oz</b>	<b>4 h</b>	<b>0 d</b>	<b>19</b>	<b>NR</b>	<b>Ph-D is similar to OSO5%SC. Use only one of these fungicides in a given season.</b>
	<b>Notes: To limit the potential for development of fungicide resistance, rotate between FRAC groups. Do not make sequential applications of fungicides in the same FRAC group. Do not make more than 2 applications of fungicides in the same FRAC group per season.</b>						
<b>Botrytis-Gray Mold</b>	<b>fluxapyroxad + pyraclostrobin Merivon</b>	<b>5.5 fl oz</b>	<b>12 h</b>	<b>0 d</b>	<b>7, 11</b>	<b>3</b>	<b>Merivon is a FRAC group 7 fungicide and should not be rotated or used with other FRAC 7 fungicides. Do not make more than three FRAC 7 applications in a season.</b>
	<b>pyraclostrobin + boscalid Pristine</b>	<b>18.5 oz</b>	<b>12 h</b>	<b>0 d</b>	<b>11,7</b>	<b>4</b>	<b>Pristine is a FRAC 7 +11 fungicide and should not be rotated or used with other FRAC 7+11 fungicides. To limit the potential for development of fungicide resistance, do not make more than 3 applications of FRAC 7+11 fungicides per season. Do not make sequential applications of FRAC 7+11 fungicides labeled for use on pome fruits. The efficacy level will depends on the occurrence of fungicide resistant populations of the pathogen in the orchards. For powdery mildew, preferably use other FRAC 7 fungicides in spring.</b>
<b>Fire blight</b>	<b>acibenzolar-s-methyl Actigard 50WG</b>	<b>2 fl oz</b>	<b>12 h</b>	<b>60 d</b>	<b>P01</b>	<b>NR</b>	<b>For bloom applications: Apply 2 oz/A in a tank mix with a fire blight treatment (generally an antibiotic) that is standard in your area. This is generally 2-3 applications between 20% bloom and petal fall depending on the environmental conditions. Do not apply closer than a 7-day interval. Also used to reduce re-occurrence of blight after cutting out infected strikes. Apply concentrate to a one meter section of the main leader after cutting see <a href="http://treefruit.wsu.edu/crop-protection/disease-management/fire-blight/">http://treefruit.wsu.edu/crop-protection/disease-management/fire-blight/</a></b>
	<b>Aureobasidium pullulans strains DSM 14940 &amp; 14941 Blossom Protect</b>	<b>1.25 lb</b>	<b>4 h</b>	<b>none listed</b>		<b>4</b>	<b>Apply with Buffer Protect. 50% and 80% bloom. Yeasts need 1-2 days before an infection to colonize the flower before bacteria invade to be effective. <span style="border: 1px solid black; padding: 0 2px;">Organic</span></b>
	<b>copper octanoate Cueva</b>	<b>4 qt</b>	<b>4 h</b>	<b>0 d</b>	<b>M1</b>	<b>3</b>	<b>Little russet in semi-arid WA trials. Some russet risk in wetter OR. Tank mix compatible with Bacillus-based biopesticides. <span style="border: 1px solid black; padding: 0 2px;">Organic</span></b>
	<b>Bacillus amyloliquefaciens strain D747 DoubleNickel 55</b>	<b>See label</b>	<b>4 h</b>	<b>0 d</b>		<b>2</b>	<b>See label and space between rows to select the corresponding rate. Efficacy may vary based on disease pressure. Can be used with copper fungicides to increase control. <span style="border: 1px solid black; padding: 0 2px;">Organic</span></b>

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
	<b>oxytetracycline FireLine 17WP</b>	See label	12 h	60 d	41	4	Best activity within 24 h before wetness event. Check spray tank pH, 5.5-6.0 optimal. Best activity at 200 ppm: 1.0 lb/100 gal. Label allows up to 1.5 lb/A. Do not go above 150 gal/A to maintain 200 ppm.
	<b>Copper Sulfate + Pentahydrate (metallic copper 5.4%) Instill</b>	30 fl oz	48 h	0 d	M1	3	Consider drying conditions to minimize marking risk.
	<b>copper sulfate pentahydrate Instill-O</b>	51 fl oz	48 h		M1	3	Consider drying conditions to minimize marking risk. <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<b>kasugamycin Kasumin 2L</b>	64 oz	12 h	90 d	24	4	Best control when applied less than 24 hrs before wetness event. Control up to 12 hr after wetness event.
	<b>calcium oxytetracycline Mycoshield</b>	See Label	12 h	60 d	41	4	Best activity within 24 h before wetness event. Check spray tank pH, 5 optimal. 200 ppm: 1.0 lb/100 gal.
	<b>copper hydroxide Previsto</b>	3-4 qt	48 h	none listed	M1	3	Pay attention to drying times and do not combine with acidifying products to reduce fruit finish risks. <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<b>Bacillus subtilis strain QST 713 Serenade Opti</b>	20 oz	4 h	0 d	44	NR	Efficacy may vary based on disease pressure. <span style="border: 1px solid black; padding: 2px;">Organic</span>
<p><b>Notes: Full bloom to petal fall. Watch the model. If an infection event occurs, apply an antibiotic as soon as possible, but within 24 hours of infection (usually wetting of flowers). Repeated antibiotic sprays may be necessary during extended high or extreme risk periods. Best results are obtained when applied within the 24-hour window before flower wetting during a high infection risk period. Beneficial only for flower infection prevention. Product used must contact the interior of the flowers in sufficient water and approved wetting agent to completely wet the interior. One pound of any 17% oxytetracycline product per 100 gallons gives a 200 ppm solution. Kasugamycin is another effective antibiotic. Applications of less than 100 gal/A can be effective on small trees if flower interiors are well covered, but do not drop the ppm below 200 (oxytetracycline). Application by ground equipment on each row is highly recommended. Application of antibiotics by aircraft is not recommended. Many fire blight bacteria in the Pacific Northwest are resistant to streptomycin, another registered antibiotic.</b></p> <p><b>Organic: Depending on the cultivar russet risk and the CougarBlight model risk follow biological with copper hydroxide/octanoate (Cueva/Previsto) every 5 to 6 days (this option is less fruit safe for russet but has higher efficacy) or with Bacillus subtilis (Serenade Opti) (most fruit safe, lower efficacy) every 2-5 days during flower/petal fall.</b></p>							
<b>Pathogens of Storage Rots</b>	<b>penthiopyrad Fontelis</b>	20 fl oz	12 h	28 d	7	3	Fontelis has fair efficacy against Alternaria fungus and Sphaeropsis that may infect fruit preharvest. Fontelis is a FRAC 7 fungicide and should not be rotated or used with other FRAC 7 fungicides. To limit the potential for development of fungicide resistance, do not make more than 3 applications of FRAC 7 fungicides per season. Do not make sequential applications of FRAC 7 fungicides labeled for use on pome fruits.

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
	<b>polyoxin D zinc salt OSO 5%SC</b>	13 fl oz	4 h	0 d	19	3	
Insect	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
<b>Campyloomma</b>	<b>acetamiprid Assail 70WP</b>	1.7-3.4 oz	12 h	7 d	4A	NR	<i>Use higher rates for high population numbers. May be applied to blooming plant in late evening, do not begin spraying until 6 pm, and stop spraying at midnight. See Pollinator Protection section.</i>
<b>Codling moth</b>	<b>CM pheromone dispensers Isomate-C Plus</b>	See label	none listed	none listed		NR	<i>Install dispensers before first flight (prior to bloom) using the full label rate in the top 2 feet of the canopy. When using aerosol emitters borders should be treated with hand-applied dispensers. <span style="border: 1px solid black; padding: 0 2px;">Organic</span></i>
<b>Leafrollers (Pandemis, Obliquebanded)</b>	<b>Bacillus thuringiensis subsp. kurstaki DiPel DF</b>	See label	4 h	0 d	11B2	3	<i>Bts are stomach poisons so complete coverage is very important for control. Two or three applications are usually required. Apply when forecasts predict a warm weather pattern, above 65°F, for 3 or more days. <span style="border: 1px solid black; padding: 0 2px;">Organic</span></i>
	<b>methoxyfenozide Intrepid 2F</b>	16 fl oz	4 h	14 d	18A	3	<i>Some leafroller populations have developed resistance to Intrepid and in these situations, the level of control can be significantly reduced.</i>
<b>Notes: Target leafroller larvae in the spring at 300-560 DD (Obliquebanded) and 300-460 DD (Pandemis). Target 1st summer generation larvae at 955-1300 DD (Obliquebanded) and 1760-2255 DD (Pandemis).</b>							

Efficacy numbers denote the relative efficacy of a pesticide against a given pest on a 1 to 4 scale with 1 being low and 4 high efficacy. This information is based primarily on research conducted with WSU researchers in Washington.

## Petal fall

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
<b>Apple mildew</b>	<b>trifloxystrobin Flint Extra</b>	2.9 fl oz	12 h	14 d	11	3	
	<b>penthiopyrad Fontelis</b>	14-20 fl oz	12 h	28 d	7	3	<i>Fontelis is a FRAC 7 fungicide.</i>
	<b>flutianil Gatten Fungicide</b>	8 fl oz	12 h	14 d	U13	NR	<i>The mechanism of action of Gatten is yet unknown. Rotate with fungicides from different chemical groups to reduce fungicide resistance.</i>

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
	<b>cyprodinil + difenoconazole</b> <i>Inspire Super</i>	12 fl oz	12 h	14 d	9,3	3	<i>Inspire Super contains cyprodinil, an active ingredient similar to the one in Vanguard. Do not rotate this product or use in the same season to avoid fungicide resistance development. Do apply after or before Rally 40WSP or other FRAC group 3 fungicides to minimize fungicide resistance in powdery mildew and other non-target pathogens.</i>
	<b>potassium bicarbonate</b> <i>Kaligreen</i>	3 lb	4 h	1 d	NC	2	<i>Under low disease pressure, Kaligreen will provide adequate control. If disease pressure increases, rotation or combination with other materials is recommended.</i> <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<b>fluopyram</b> <i>Luna Privilege</i>	6.84 fl oz	12 h	7 d	7	4	<i>Luna Privilege is a FRAC group 7 fungicide.</i>
	<b>fluopyram + trifloxystrobin</b> <i>Luna Sensation</i>	5-5.8 fl oz	12 h	14 d	7, 11	4	
	<b>fluxapyroxad + pyraclostrobin</b> <i>Merivon</i>	5.5 fl oz	12 h	0 d	7, 11	3	<i>Merivon is a FRAC group 7 fungicide.</i>
	<b>polyoxin D zinc salt</b> <i>OSO 5%SC</i>	13 fl oz	4 h	0 d	19	3	<i>Do not apply more than 4.2 oz. a.i./acre/season.</i>
	<b>polyoxin D zinc salt</b> <i>Ph-D</i>	6.2 oz	4 h	0 d	19	NR	<i>Ph-D is similar to OSO5%SC. Use only one of these fungicides in a given season.</i>
	<b>pyraclostrobin + boscalid</b> <i>Pristine</i>	18.5 oz	12 h	0 d	11,7	3	<i>Pristine is a FRAC 7 +11 fungicide and should not be rotated or used with other FRAC 7+11 fungicides. The efficacy level will depend on the occurrence of fungicide resistant populations of the pathogen in the orchards. For powdery mildew, preferably use other FRAC 7 fungicides in spring.</i>
	<b>myclobutanil</b> <i>Rally 40WSP</i>	5 oz	24 h	14 d	3	4	<i>Rally is a FRAC 3 fungicide and should not be rotated or used with other FRAC 3 fungicides. Place Rally into solution before adding oil. Apply no sooner than half-inch green.</i>
	<b>Reynoutria sachalinensis</b> <i>Regalia</i>	4 qt	4 h	0 d	P5	2	<i>Do not use prior to petal fall. Under low disease pressure, it may help control some summer diseases like Alternaria and Bull's eye rot.</i> <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<b>lime sulfur/calcium polysulfide</b> <i>Rex Lime Sulfur</i>	See label	48 h	0 d		NR	<i>Do not apply lime sulfur if temperatures will exceed 75°F within 3 days of application.</i> <span style="border: 1px solid black; padding: 2px;">Organic</span>

<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<b>Bacillus pumilus strain QST 2808 Sonata</b>	2-4 qt	4 h	0 d		NR	Organic
	<b>flutriafol Topguard</b>	8-12 fl oz	12 h	14 d	3	4	Topguard is a FRAC group 3 fungicide and should not be rotated or used with other FRAC 3 fungicides. Topguard may cause a mild phytotoxicity on Braeburn apple leaves.
	<b>triflumazole Trionic 4SC</b>	16 fl oz	12 h	14 d	3	4	Trionic is a FRAC 3 fungicide and should not be rotated or used with other FRAC 3 fungicides.
	<b>Notes: The efficacy level will depend on the absence of resistant populations from the orchard. To limit the potential for development of fungicide resistance, do not make more than four applications of FRAC group 11 fungicides per season. Do not make more than two sequential applications of FRAC group 11 fungicides. This limitation is inclusive of all FRAC group 11 fungicides labeled for use on pome fruits.</b>						
	<b>Notes: To limit the potential for development of fungicide resistance, rotate between FRAC groups. Do not make sequential applications of fungicides in the same FRAC group. Do not make more than 2 applications of fungicides in the same FRAC group per season.</b>						
<b>Apple scab</b>	<b>captan Captan 50WP</b>	6 lb	24 h	0 d	M4	NR	When possible, tank-mix captan with other single-site fungicides to help reduce fungicide resistance development.
	<b>trifloxystrobin Flint Extra</b>	2.5-2.9 fl oz	12 h	14 d	11	3	
	<b>penthiopyrad Fontelis</b>	14-20 fl oz	12 h	28 d	7	NR	
	<b>fluopyram + trifloxystrobin Luna Sensation</b>	4-5.8 fl oz	12 h	14 d	7, 11	NR	Luna Sensation, Fontelis, and Aprovia are all from the same chemical group (7). Use one of them ONLY.
	<b>metiram Polyram 80DF</b>	See label	24 h	77 d	M3	NR	
	<b>triflumazole Procure 480SC</b>	8-16 fl oz	12 h	14 d	3	NR	
	<b>myclobutanil Rally 40WSP</b>	5 oz	24 h	14 d	3	NR	Use only one Group 3 fungicide at the same growth stage
	<b>flutriafol Topguard</b>	13 fl oz	12 h	14 d	3	NR	Topguard may cause a mild phytotoxicity on Braeburn apple leaves. Use only one Group 3-fungicide at the same growth stage.
	<b>triflumazole Trionic 4SC</b>	12-16 fl oz	12 h	14 d	3	NR	Use only one Group-3 fungicide at the same growth stage.

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
	ziram Ziram 76DF	See label	48 h	14 d	M3	NR	
<p><b>Notes: To limit the potential for development of fungicide resistance, do not make more than four applications of strobilurin fungicides per season. Do not make more than two sequential applications of strobilurin fungicides. This limitation is inclusive of all strobilurin fungicides (Mode 11) labeled for use on pome fruits.</b></p>							
<b>Botrytis-Gray Mold</b>	captan Captan 50WP	8 lb	24 h	0 d	M4	2	Do not apply more than 64 lbs. of Captan 50 Wettable Powder per acre per crop cycle.
	copper octanoate Cueva	8 qt	4 h	0 d	M1	2	Organic
	fluxapyroxad + pyraclostrobin Merivon	5.5 fl oz	12 h	0 d	7, 11	3	Merivon is a FRAC group 7 fungicide and should not be rotated or used with other FRAC 7 fungicides. Do not make more than three FRAC 7 applications in a season.
	pyraclostrobin + boscalid Pristine	18.5 oz	12 h	0 d	11,7	4	Pristine is a FRAC 7 +11 fungicide and should not be rotated or used with other FRAC 7+11 fungicides. To limit the potential for development of fungicide resistance, do not make more than 3 applications of FRAC 7+11 fungicides per season. Do not make sequential applications of FRAC 7+11 fungicides labeled for use on pome fruits. The efficacy level will depends on the occurrence of fungicide resistant populations of the pathogen in the orchards. For powdery mildew, preferably use other FRAC 7 fungicides in spring.
	thiophanate-methyl Topsin M WSB	1 lb	2 d	1 d	1	3	Efficacy level is ensured when resistance is absent from the orchard. Do not apply more than 2.8 lbs. a.i. per acre in a year. Topsin-M is very prone to fungicide resistance development, rotation with other FRAC group is required. Do not make sequential applications of FRAC group 1 fungicides and do not make more than two FRAC 1 fungicides per season. Topsin-M is similar to Mertect, used postharvest, therefore careful use is highly recommended.

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
Fire blight	acibenzolar-s-methyl Actigard 50WG	2 fl oz	12 h	60 d	P01	NR	For bloom applications: Apply 2 oz/A in a tank mix with a fire blight treatment (generally an antibiotic) that is standard in your area. This is generally 2-3 applications between 20% bloom and petal fall depending on the environmental conditions. Do not apply closer than a 7-day interval. Also used to reduce re-occurrence of blight after cutting out infected strikes. Apply concentrate to a one meter section of the main leader after cutting see <a href="http://treefruit.wsu.edu/crop-protection/disease-management/fire-blight/">http://treefruit.wsu.edu/crop-protection/disease-management/fire-blight/</a>
	copper octanoate Cueva	4 qt	4 h	0 d	M1	3	Little russet in semi-arid WA trials. Some russet risk in wetter OR. Tank mix compatible with Bacillus-based biopesticides. <span>Organic</span>
	Bacillus amyloliquifaciens strain D747 DoubleNickel 55	See label	4 h	0 d		2	See label and space between rows to select the corresponding rate. Efficacy may vary based on disease pressure. Can be used with copper fungicides to increase control. <span>Organic</span>
	oxytetracycline FireLine 17WP	See label	12 h	60 d	41	4	Best activity within 24 h before wetness event. Check spray tank pH, 5.5-6.0 optimal. Best activity at 200 ppm: 1.0 lb/100 gal. Label allows up to 1.5 lb/A. Do not go above 150 gal/A to maintain 200 ppm.
	kasugamycin Kasumin 2L	64 oz	12 h	90 d	24	4	Best control when applied less than 24 hrs before wetness event. Control up to 12 hr after wetness event.
	calcium oxytetracycline Mycoshield	See Label	12 h	60 d	41	4	Best activity within 24 h before wetness event. Check spray tank pH, 5 optimal. 200 ppm: 1.0 lb/100 gal.
	lime sulfur/calcium polysulfide lime sulfur/calcium polysulfide	2-4 % v/v	48 h	none listed		NR	At petal fall lime sulfur (2 to 4%) to clean up bacteria, yeast, mildew and rot fungi.
	copper hydroxide Previsto	3-4 qt	48 h	none listed	M1	3	Pay attention to drying times and do not combine with acidifying products to reduce fruit finish risks. <span>Organic</span>
	Bacillus subtilis strain QST 713 Serenade Opti	20 oz	4 h	0 d	44	NR	Efficacy may vary based on disease pressure. <span>Organic</span>
	Pathogens of Storage Rots	Bacillus subtilis strain IAB/BS03 Aviv	10-30 fl oz	4 h	none listed	44	3

<i>Disease</i>	<i>Chemical</i>	<i>Rate per Acre</i>	<i>REI</i>	<i>PHI</i>	<i>MOA</i>	<i>Eff.</i>	<i>Notes</i>
	<b>penthiopyrad</b> <b>Fontelis</b>	20 fl oz	12 h	28 d	7	3	<i>Fontelis has fair efficacy against Alternaria fungus and Sphaeropsis that may infect fruit preharvest. Fontelis is a FRAC 7 fungicide and should not be rotated or used with other FRAC 7 fungicides. To limit the potential for development of fungicide resistance, do not make more than 3 applications of FRAC 7 fungicides per season. Do not make sequential applications of FRAC 7 fungicides labeled for use on pome fruits.</i>
	<b>polyoxin D zinc salt</b> <b>OSO 5%SC</b>	13 fl oz	4 h	0 d	19	3	<i>OSO will help control Alternaria infections preharvest. OSO is FRAC 19 fungicide recommended to rotate with other FRAC groups labeled for pome fruit. Do not apply more than 4.2 oz. a.i./acre/season.</i>
	<b>polyoxin D zinc salt</b> <b>Ph-D</b>	6.2 oz	4 h	0 d	19	NR	<i>Ph-D is similar to OSO 5%, so use only one of them in a given season.</i>
<i>Insect</i>	<i>Chemical</i>	<i>Rate per Acre</i>	<i>REI</i>	<i>PHI</i>	<i>MOA</i>	<i>Eff.</i>	<i>Notes</i>
<b>Codling moth</b>	<b>chlorantraniliprole</b> <b>Altacor</b>	4.5 oz	4 h	5 d	28	4	<i>Altacor is highly effective against leafroller larvae and, at this treatment timing, has added value because it is toxic to codling moth eggs laid on product residues (see recommendations under codling moth). Altacor can, therefore, be used as part of a management strategy to delay the first larvicide application against codling moth. Use the leafroller models on the WSU Decision Aid System (<a href="https://decisionaid.systems">https://decisionaid.systems</a>) for the optimum timing for this product.</i>
	<b>spinetoram</b> <b>Delegate WG</b>	7 oz	4 h	7 d	5	4	<i>Delegate is highly effective against leafroller larvae. While Delegate does not directly kill codling moth eggs it has a strong ovi-larvicidal activity, which means it kills codling moth larvae exiting eggs. Therefore, if Delegate is applied at this timing it can be used as part of a management strategy to delay the first larvicide application against codling moth. Delegate is in the same chemical class, and has the same mode of action, as Success (spinosad) so avoid using these products against two consecutive generations.</i>
	<b>pyriproxyfen</b> <b>Esteem 35WP</b>	4-5 oz	12 h	45 d	7C	3	<i>Should be applied when the last larval stage is present but before pupation has begun. Use the WSU Decision Aid System (<a href="https://decisionaid.systems">https://decisionaid.systems</a>) for the optimum timing of the product on leafrollers.</i>



<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<i>methoxyfenozide</i> <b>Intrepid 2F</b>	16 fl oz	4 h	14 d	18A	3	
	<i>novaluron</i> <b>Rimon 0.83EC</b>	30-50 fl oz	12 h	14 d	15	3-4	
	<i>petroleum oil, summer</i> <i>petroleum oil, summer</i>	See label	4 h	0 d		NR	Organic
<p><b>Notes: WSU recommends a delayed first cover approach. Apply the first oil at 375 DD, then 150 degree days later put on the first cover at 525 DD. Then 15 days later (depending on residue length) put on the second cover. This approach leaves only a small percentage of egg hatch at the end of each generation uncovered. See <a href="https://decisionaid.systems">https://decisionaid.systems</a></b></p>							
<b>Grape mealybug</b>	<i>imidacloprid</i> <b>Admire Pro</b>	7 fl oz	12 h	7 d	4A	NR	Rate/PHI for foliar application.
	<i>acetamiprid</i> <b>Assail 70WP</b>	3.4 oz	12 h	7 d	4A	1	
	<i>buprofezin</i> <b>Centaur WDG</b>	34.5 oz	12 h	14 d	16	NR	
<b>Leafrollers (Pandemis, Obliquebanded)</b>	<i>chlorantraniliprole</i> <b>Altacor</b>	3-4.5 oz	4 h	5 d	28	4	<b>Altacor is highly effective against leafroller larvae and, at this treatment timing, has added value because it is toxic to codling moth eggs laid on product residues (see recommendations under codling moth). Altacor can, therefore, be used as part of a management strategy to delay the first larvicide application against codling moth. Use the leafroller models on the WSU Decision Aid System (<a href="https://decisionaid.systems">https://decisionaid.systems</a>) for the optimum timing for this product.</b>
	<i>spinetoram</i> <b>Delegate WG</b>	4.5-7 oz	4 h	7 d	5	4	<b>Delegate is highly effective against leafroller larvae. While Delegate does not directly kill codling moth eggs it has a strong ovi-larvicidal activity, which means it kills codling moth larvae exiting eggs. Therefore, if Delegate is applied at this timing is can be used as part of a management strategy to delay the first larvicide application against codling moth. Delegate is in the same chemical class, and has the same mode of action, as Success (spinosad) so avoid using these products against two consecutive generations.</b>
	<i>Bacillus thuringiensis subsp. kurstaki</i> <b>DiPel DF</b>	See label	4 h	0 d	11B2	3	<b>Bts are stomach poisons so complete coverage is very important for control. Two or three applications are usually required. Apply when forecasts predict a warm weather pattern, above 65°F, for 3 or more days.</b> Organic

<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<b>spinosad Entrust SC</b>	<b>10 fl oz</b>	<b>4 h</b>	<b>7 d</b>	<b>5</b>	<b>NR</b>	<b>Entrust is a spinosad formulation registered for organic apple production. <span style="border: 1px solid black; padding: 2px;">Organic</span></b>
	<b>pyriproxyfen Esteem 35WP</b>	<b>4-5 oz</b>	<b>12 h</b>	<b>45 d</b>	<b>7C</b>	<b>4</b>	<b>Should be applied when the last larval stage is present but before pupation has begun. Use the WSU Decision Aid System (<a href="https://decisionaid.systems">https://decisionaid.systems</a>) for the optimum timing of the product on leafrollers.</b>
	<b>cyantraniliprole Exirel</b>	<b>10-17 fl oz</b>	<b>12 h</b>	<b>3 d</b>	<b>28</b>	<b>4</b>	
	<b>methoxyfenozide Intrepid 2F</b>	<b>16 fl oz</b>	<b>4 h</b>	<b>14 d</b>	<b>18A</b>	<b>3</b>	<b>Some leafroller populations have developed resistance to Intrepid and in these situations, the level of control can be significantly reduced.</b>
	<b>emamectin benzoate Proclaim</b>	<b>3.2-4.8 oz</b>	<b>12 h / 48 h for some activities-see label</b>	<b>14 d</b>	<b>6</b>	<b>4</b>	
	<b>novaluron Rimon 0.83EC</b>	<b>30-50 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>15</b>	<b>3-4</b>	
	<b>spinosad Success</b>	<b>6-10 fl oz</b>	<b>4 h</b>	<b>7 d</b>	<b>5</b>	<b>3-4</b>	<b>Some leafroller populations have developed resistance to Success so use of this product in this situation may result in reduced control.</b>
<b>Notes: Target leafroller larvae in the spring at 300-560 DD (Obliquebanded) and 300-460 DD (Pandemis). Target 1st summer generation larvae at 955-1300 DD (Obliquebanded) and 1760-2255 DD (Pandemis).</b>							
<b>Rosy apple aphid</b>	<b>imidacloprid Admire Pro</b>	<b>1.4 - 2.8 fl oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4A</b>	<b>NR</b>	
	<b>acetamiprid Assail 70WP</b>	<b>1.7 oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4A</b>	<b>3-4</b>	
<b>Notes: Use of these materials at this timing will also control other aphid species.</b>							
<b>Spider mites</b>	<b>bifenazate Acramite 50WS</b>	<b>0.75-1 lb</b>	<b>12 h</b>	<b>7 d</b>	<b>un</b>	<b>4</b>	
	<b>clofentezine Apollo 4SC</b>	<b>6-8 fl oz</b>	<b>12 h</b>	<b>45 d</b>	<b>10A</b>	<b>NR</b>	
	<b>spirodiclofen Envidor 2SC</b>	<b>16-18 fl oz</b>	<b>12 h</b>	<b>7 d</b>	<b>23</b>	<b>3-4</b>	
	<b>fenpyroximate FujiMite XLO</b>	<b>1-2 pt</b>	<b>12 h</b>	<b>14 d</b>	<b>21A</b>	<b>3-4</b>	

<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<b>pyridaben Nexter 75WSB</b>	<b>4.4-8.8 oz</b>	<b>12 h</b>	<b>25 d</b>	<b>21A</b>	<b>2-3</b>	
	<b>hexythiazox Savey 50DF</b>	<b>4-6 oz</b>	<b>12 h</b>	<b>28 d</b>	<b>10A</b>	<b>NR</b>	<b>Most effective on the egg stage. When mite populations are high and leaf bronzing has already occurred, a miticide effective on the adult stage may be used in combination.</b>
	<b>etoxazole Zeal Miticide1 72WSP</b>	<b>2-3 oz</b>	<b>12 h</b>	<b>14 d</b>	<b>10B</b>	<b>3-4</b>	
<b>Western flower thrips</b>	<b>spinosad Entrust SC</b>	<b>6-8 fl oz</b>	<b>4 h</b>	<b>7 d</b>	<b>5</b>	<b>3</b>	<b>Organic</b>
	<b>spinosad Success</b>	<b>6-8 fl oz</b>	<b>4 h</b>	<b>7 d</b>	<b>5</b>	<b>3</b>	
	<b>Notes: Timing tests for minimizing thrips damage indicate petal fall (5 mm fruit) is better than the traditional bloom timing.</b>						
<b>Western tentiform leafminer</b>	<b>abamectin Agri-Mek SC</b>	<b>4.25 fl oz</b>	<b>12 h</b>	<b>28 d</b>	<b>6</b>	<b>4</b>	
	<b>spinosad Success</b>	<b>6 fl oz</b>	<b>4 h</b>	<b>7 d</b>	<b>5</b>	<b>3-4</b>	
<b>White apple leafhopper</b>	<b>imidacloprid Admire Pro</b>	<b>1.4-2.8 fl oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4A</b>	<b>4</b>	<b>Do not use until pollination is complete and bees have been removed from the area. Rate indicated is for foliar application.</b>
	<b>indoxacarb Avaunt</b>	<b>6 oz</b>	<b>12 h</b>	<b>14 d</b>	<b>22</b>	<b>3-4</b>	
	<b>spinosad Success</b>	<b>6-8 fl oz</b>	<b>4 h</b>	<b>7 d</b>	<b>5</b>	<b>3</b>	
	<b>kaolin clay Surround WP</b>	<b>See label</b>	<b>4 h</b>	<b>0 d</b>		<b>NR</b>	<b>Organic</b>
	<b>Notes: Do not use until pollination is complete and bees have been removed from the area.</b>						

Efficacy numbers denote the relative efficacy of a pesticide against a given pest on a 1 to 4 scale with 1 being low and 4 high efficacy. This information is based primarily on research conducted with WSU researchers in Washington.

14-28 days after full bloom

<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
<b>Apple mildew</b>	<b>penthiopyrad Fontelis</b>	<b>14-20 fl oz</b>	<b>12 h</b>	<b>28 d</b>	<b>7</b>	<b>3</b>	<b>Fontelis is a FRAC 7 fungicide.</b>
	<b>flutianil Gatten Fungicide</b>	<b>8 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>U13</b>	<b>NR</b>	<b>The mechanism of action of Gatten is yet unknown. Rotate with fungicides from different chemical groups to reduce fungicide resistance.</b>
	<b>cyprodinil + difenoconazole Inspire Super</b>	<b>12 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>9,3</b>	<b>3</b>	<b>Inspire Super contains cyprodinil, an active ingredient similar to the one in Vanguard. Do not rotate this product or use in the same season to avoid fungicide resistance development. Do apply after or before Rally 40WSP or other FRAC group 3 fungicides to minimize fungicide resistance in powdery mildew and other non-target pathogens.</b>
	<b>potassium bicarbonate Kaligreen</b>	<b>3 lb</b>	<b>4 h</b>	<b>1 d</b>	<b>NC</b>	<b>2</b>	<b>Under low disease pressure, Kaligreen will provide adequate control. If disease pressure increases, rotation or combination with other materials is recommended.</b> <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<b>fluopyram Luna Privilege</b>	<b>6.84 fl oz</b>	<b>12 h</b>	<b>7 d</b>	<b>7</b>	<b>4</b>	<b>Luna Privilege is a FRAC group 7 fungicide.</b>
	<b>fluxapyroxad + pyraclostrobin Merivon</b>	<b>5.5 fl oz</b>	<b>12 h</b>	<b>0 d</b>	<b>7, 11</b>	<b>3</b>	<b>Merivon is a FRAC group 7 fungicide.</b>
	<b>polyoxin D zinc salt OSO 5%SC</b>	<b>13 fl oz</b>	<b>4 h</b>	<b>0 d</b>	<b>19</b>	<b>3</b>	<b>Do not apply more than 4.2 oz. a.i./acre/season.</b>
	<b>polyoxin D zinc salt Ph-D</b>	<b>6.2 oz</b>	<b>4 h</b>	<b>0 d</b>	<b>19</b>	<b>NR</b>	<b>Ph-D is similar to OSO5%SC. Use only one of these fungicides in a given season.</b>
	<b>pyraclostrobin + boscalid Pristine</b>	<b>18.5 oz</b>	<b>12 h</b>	<b>0 d</b>	<b>11,7</b>	<b>3</b>	<b>Pristine is a FRAC 7 +11 fungicide and should not be rotated or used with other FRAC 7+11 fungicides. The efficacy level will depend on the occurrence of fungicide resistant populations of the pathogen in the orchards. For powdery mildew, preferably use other FRAC 7 fungicides in spring.</b>
	<b>triflumizole Procure 480SC</b>	<b>8-16 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>3</b>	<b>4</b>	
	<b>myclobutanil Rally 40WSP</b>	<b>5 oz</b>	<b>24 h</b>	<b>14 d</b>	<b>3</b>	<b>4</b>	<b>Rally is a FRAC 3 fungicide and should not be rotated or used with other FRAC 3 fungicides. Place Rally into solution before adding oil. Apply no sooner than half-inch green.</b>

<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<i>Reynoutria sachalinensis</i> Regalia	4 qt	4 h	0 d	P5	2	Do not use prior to petal fall. Under low disease pressure, it may help control some summer diseases like Alternaria and Bull's eye rot. <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<i>Bacillus pumilus</i> strain QST 2808 Sonata	2-4 qt	4 h	0 d		NR	<span style="border: 1px solid black; padding: 2px;">Organic</span>
	flutriafol Topguard	8-12 fl oz	12 h	14 d	3	4	Topguard is a FRAC group 3 fungicide and should not be rotated or used with other FRAC 3 fungicides. Topguard may cause a mild phytotoxicity on Braeburn apple leaves.
<p><b>Notes: The efficacy level will depend on the absence of propiconazole-resistant populations from the orchard. To limit the potential for development of fungicide resistance, do not make more than four applications of FRAC group 11 fungicides per season. Do not make more than two sequential applications of FRAC group 11 fungicides. This limitation is inclusive of all FRAC group 11 fungicides labeled for use on pome fruits.</b></p>							
<b>Apple scab</b>	captan Captan 50WP	6 lb	24 h	0 d	M4	NR	
	penthiopyrad Fontelis	14-20 fl oz	12 h	28 d	7	NR	Fontelis and Luna Sensation are all from the same chemical group (7). Apply ONLY one of them at the same growth stage.
	cyprodinil + difenoconazole Inspire Super	12 fl oz	12 h	14 d	9,3	3	Inspire Super contains cyprodinil, an active ingredient similar to the one in Vanguard. Do not rotate this products or use in the same season to avoid fungicide resistance development. Do apply after or before Rally 40WSP or other FRAC group 3 fungicides to minimize fungicide resistance in powdery mildew and other non-target pathogens.
	fluopyram + trifloxystrobin Luna Sensation	4-5.8 fl oz	12 h	14 d	7, 11	NR	Luna Sensation and Fontelis are from the same Chemical group (7). Use only one of them.
	metiram Polyram 80DF	See label	24 h	77 d	M3	NR	
	triflumizole Procure 480SC	8-16 fl oz	12 h	14 d	3	NR	Rates vary with postinfective schedule; see label. Apply only one Group 3 Fungicide at the same growth stage.
	myclobutanil Rally 40WSP	5 oz	24 h	14 d	3	NR	
	flutriafol Topguard	13 fl oz	12 h	14 d	3	NR	Topguard may cause a mild phytotoxicity on Braeburn apple leaves.

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
	ziram Ziram 76DF	See label	48 h	14 d	M3	NR	
<p><b>Notes: To limit the potential for development of fungicide resistance, do not make more than four applications of strobilurin fungicides per season. Do not make more than two sequential applications of strobilurin fungicides. This limitation is inclusive of all strobilurin fungicides (Mode 11) labeled for use on pome fruits.</b></p>							
<b>Botrytis-Gray Mold</b>	pyraclostrobin + boscalid Pristine	18.5 oz	12 h	0 d	11,7	4	Pristine is a FRAC 7 +11 fungicide and should not be rotated or used with other FRAC 7+11 fungicides. To limit the potential for development of fungicide resistance, do not make more than 3 applications of FRAC 7+11 fungicides per season. Do not make sequential applications of FRAC 7+11 fungicides labeled for use on pome fruits. The efficacy level will depends on the occurrence of fungicide resistant populations of the pathogen in the orchards. For powdery mildew, preferably use other FRAC 7 fungicides in spring.
<b>Bull's eye rot</b>	ziram Ziram Granuflo 76WDG	6 lb	48 h	14 d	M3	3	Apply 12 to 14 days after calyx stage and repeat as needed through the summer. Do not apply more than 24.2 lbs per acre in a crop cycle. Do not apply within 14 days of harvest. Tank-mixing with other fungicides has been reported to increase efficacy and reduce fungicide resistance development.
<b>Pathogens of Storage Rots</b>	polyoxin D zinc salt OSO 5%SC	13 fl oz	4 h	0 d	19	3	OSO will help control Alternaria infections preharvest. OSO is FRAC 19 fungicide recommended to rotate with other FRAC groups labeled for pome fruit. Do not apply more than 4.2 oz. a.i./acre/season.
	polyoxin D zinc salt Ph-D	6.2 oz	4 h	0 d	19	NR	Ph-D is similar to OSO 5%, so use only one of them in a given season.
Insect	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
<b>Codling moth</b>	chlorantraniliprole Altacor	3-4.5 oz	4 h	5 d	28	4	Altacor is effective against codling moth eggs and larvae. At the high rate, it has a residual activity of up to 17 days, at the low rate residual activity is no more than 14 days.
	acetamiprid Assail 70WP	3.4 oz	12 h	7 d	4A	NR	If applied to control codling moth, this product will provide control of rosy apple aphid as well at this timing. Use an appropriate surfactant to enhance coverage and penetration.

<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<b>CM granulosis virus (CpGV) Cyd-X</b>	<b>See Label</b>	<b>4 h</b>	<b>0 d</b>		<b>3</b>	<b>Codling moth granulosis virus is a highly specific control that should always be used as a component of a multi-tactic pest management program. The residual activity lasts at most 7 days. The effect of the virus is most often seen in a suppression of the pest's densities over time. Applying virus one or more times per codling moth generation at the end of the residual period of another insecticide to extend the protection period is a logical use of this technology.</b> <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<b>spinetoram Delegate WG</b>	<b>6-7 oz</b>	<b>4 h</b>	<b>7 d</b>	<b>5</b>	<b>4</b>	<b>Delegate is very effective against codling moth larvae. It has a residual activity of 14 days.</b>
	<b>spinosad Entrust SC</b>	<b>3 fl oz</b>	<b>4 h</b>	<b>7 d</b>	<b>5</b>	<b>NR</b>	<b>Entrust is a spinosad formulation registered for organic apple production. It is effective against codling moth larvae hatching from the egg. It has a residual activity of 7 to 10 days. When Entrust is incorporated into an organic pest control program using pheromones, summer oil, and codling moth virus, good control of this key pest is possible. Best results occur when applications are timed for egg hatch, which may occur during bloom.</b> <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<b>cyantraniliprole Exirel</b>	<b>10-17 fl oz</b>	<b>12 h</b>	<b>3 d</b>	<b>28</b>	<b>4</b>	
	<b>phosmet Imidan 70W</b>	<b>5.33 lb</b>	<b>7 d</b>	<b>7 d</b>	<b>1B</b>	<b>3</b>	
	<b>methoxyfenozide Intrepid 2F</b>	<b>16 fl oz</b>	<b>4 h</b>	<b>14 d</b>	<b>18A</b>	<b>3</b>	<b>Intrepid is a stomach poison so complete coverage is important to good control. This product is recommended only as a supplement to mating disruption. Apply the first application of Intrepid at 425 degree days using the new codling moth model (525 DD when using the delayed first cover with oil at 375 DD) and follow with additional applications at 14-day intervals for a total of 3 sprays. Intrepid can also be used in the second generation (if not used in the first generation) timed at egg hatch and using the same re-treatment intervals.</b>
	<b>petroleum oil, summer petroleum oil, summer</b>	<b>See label</b>	<b>4 h</b>	<b>0 d</b>		<b>NR</b>	<span style="border: 1px solid black; padding: 2px;">Organic</span>
<b>Notes: For a delayed first cover program, the first larvicide should be applied at 525 degree days. For a standard (no ovicide) program, the timing is 425 degree days (see <a href="https://decisionaid.systems">https://decisionaid.systems</a>). If necessary, a second larvicide for the first generation should be applied 14-17 days after the first application.</b>							

<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
<b>Rosy apple aphid</b>	<b>imidacloprid Admire Pro</b>	<b>1.4-2.8 fl oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4A</b>	<b>NR</b>	
	<b>acetamiprid Assail 70WP</b>	<b>1.7 oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4A</b>	<b>3-4</b>	<b>If applied to control codling moth, this product will provide control of rosy apple aphid as well at this timing. Use an appropriate surfactant to enhance coverage and penetration.</b>
<b>White apple leafhopper</b>	<b>indoxacarb Avaunt</b>	<b>6 oz</b>	<b>12 h</b>	<b>14 d</b>	<b>22</b>	<b>3-4</b>	
	<b>kaolin clay Surround WP</b>	<b>See label</b>	<b>4 h</b>	<b>0 d</b>		<b>NR</b>	<b>Organic</b>
<b>Woolly apple aphid</b>	<b>sulfoxaflor Closer SC</b>	<b>4.25-5.75 fl oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4C</b>	<b>3</b>	<b>Use an adjuvant such as Exit at 0.25% vol:vol.</b>
	<b>spirotetramat Ultror</b>	<b>10-14 fl oz</b>	<b>24 h</b>	<b>7 d</b>	<b>23</b>	<b>2-4</b>	<b>Time Ultror applications shortly after petal fall, when the canopy is well developed, but leaves have not hardened off. A second application 14 days after the first may be helpful. This material is systemic and will suppress root and shoot colonies of woolly apple aphid. Ultror suppresses woolly apple aphids later in the season, but overall control is not as good as with the early season timing (see Late Spring and Summer).</b>

Efficacy numbers denote the relative efficacy of a pesticide against a given pest on a 1 to 4 scale with 1 being low and 4 high efficacy. This information is based primarily on research conducted with WSU researchers in Washington.

## Late spring and summer

<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
<b>Apple maggot</b>	<b>thiamethoxam Actara</b>	<b>5.5 oz</b>	<b>12 h</b>	<b>14 d/35 d</b>	<b>4A</b>	<b>NR</b>	<b>Estimated residual activity: 7-14 days.</b>
	<b>imidacloprid Admire Pro</b>	<b>2.8 fl oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4A</b>	<b>NR</b>	<b>Estimated residual activity: 10-14 days. Rate/PHI for foliar application.</b>
	<b>chlorantraniliprole Altacor</b>	<b>4.5 oz</b>	<b>4 h</b>	<b>5 d</b>	<b>28</b>	<b>NR</b>	<b>Provides suppression only. Also effective against codling moth and leafrollers. Estimated residual activity: 10-14 days.</b>
	<b>acetamiprid Assail 70WP</b>	<b>3.4 oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4A</b>	<b>NR</b>	<b>Assail is also used to control codling moth. The estimated residual activity is 10-14 days.</b>



<i>Insect</i>	<i>Chemical</i>	<i>Rate per Acre</i>	<i>REI</i>	<i>PHI</i>	<i>MOA</i>	<i>Eff.</i>	<i>Notes</i>
	<i>fenpropathrin</i> <i>Danitol 2.4EC</i>	16-21.3 fl oz	24 h	14 d	3	NR	<i>Also effective against codling moth and leafrollers. Estimated residual activity: 7-10 days.</i>
	<i>spinetoram</i> <i>Delegate WG</i>	6-7 oz	4 h	7 d	5	NR	<i>Provides suppression only. Also effective against codling moth and leafrollers. Estimated residual activity: 10-14 days.</i>
	<i>spinosad</i> <i>Entrust</i>	2-3 oz	4 h	7 d	5	NR	<i>Provides suppression only. Also effective against codling moth and leafrollers. Estimated residual activity: 5-7 days. <span style="border: 1px solid black; padding: 0 2px;">Organic</span></i>
	<i>phosmet</i> <i>Imidan 70W</i>	5.33 lb	7 d	7 d	1B	NR	<i>Imidan is also used to control codling moth. The estimated residual activity is 14 days.</i>
	<i>lambda-cyhalothrin</i> <i>Warrior II</i>	1.28-2.56 fl oz	24 h	21 d	3	NR	<i>Also effective against codling moth and leafrollers. Estimated residual activity: 7-10 days.</i>
<b>Codling moth</b>	<i>chlorantraniliprole</i> <i>Altacor</i>	3-4.5 oz	4 h	5 d	28	4	<i>Altacor is very effective against leafroller larvae and codling moth eggs and larvae. It has a residual activity of up to 17 days when using the high label rate or not more than 14 days when using the low label rate.</i>
	<i>acetamiprid</i> <i>Assail 70WP</i>	3.4 oz	12 h	7 d	4A	NR	
	<i>CM granulosis virus (CpGV)</i> <i>Cyd-X</i>	See Label	4 h	0 d		3	<i>Codling moth granulosis virus is a highly specific control that should always be used as a component of a multi-tactic pest management program. The residual activity lasts at most 7 days. The effect of the virus is most often seen in a suppression of the pest's densities over time. Applying virus one or more times per codling moth generation at the end of the residual period of another insecticide to extend the protection period is a logical use of this technology. <span style="border: 1px solid black; padding: 0 2px;">Organic</span></i>
	<i>spinetoram</i> <i>Delegate WG</i>	6-7 oz	4 h	7 d	5	4	<i>Delegate is very effective against leafroller and codling moth larvae. It has a residual activity of 14 days. Delegate is in the same chemical class as Success (spinosad) so avoid using these products against two consecutive generations.</i>

<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<b>spinosad Entrust SC</b>	<b>3 fl oz</b>	<b>4 h</b>	<b>7 d</b>	<b>5</b>	<b>NR</b>	<b>Entrust is a spinosad formulation registered for organic apple production. It is effective against codling moth larvae hatching from the egg. It has a residual activity of 7 to 10 days. When Entrust is incorporated into an organic pest control program using pheromones, summer oil, and codling moth virus, good control of this key pest is possible. Best results occur when applications are timed for egg hatch, which may occur during bloom.</b> <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<b>cyantraniliprole Exirel</b>	<b>10-17 fl oz</b>	<b>12 h</b>	<b>3 d</b>	<b>28</b>	<b>4</b>	
	<b>phosmet Imidan 70W</b>	<b>5 lb</b>	<b>7 d</b>	<b>7 d</b>	<b>1B</b>	<b>3</b>	
	<b>methoxyfenozide Intrepid 2F</b>	<b>16 fl oz</b>	<b>4 h</b>	<b>14 d</b>	<b>18A</b>	<b>3</b>	
	<b>novaluron Rimon 0.83EC</b>	<b>30-50 fl oz</b>	<b>12 h</b>	<b>14 d</b>	<b>15</b>	<b>3-4</b>	
	<b>petroleum oil, summer petroleum oil, summer</b>	<b>See label</b>	<b>4 h</b>	<b>0 d</b>		<b>NR</b>	<span style="border: 1px solid black; padding: 2px;">Organic</span>
	<b>Notes: If a residual ovicide is used against the second generation, it should be applied at 1375 degree days (the delayed first cover timing), and the first larvicide delayed until 1525 degree days. The standard program larvicide is applied at 1425 degree days (when no oil is used at 1375). If necessary, a second larvicide should be applied 14-17 days after the first. Timings for the third generation should add 1,000 degree days to those of the second generation.</b>						
<b>Cutworms</b>	<b>indoxacarb Avaunt</b>	<b>6 oz</b>	<b>12 h</b>	<b>14 d</b>	<b>22</b>	<b>4</b>	
<b>Grape mealybug</b>	<b>imidacloprid Admire Pro</b>	<b>7 fl oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4A</b>	<b>NR</b>	<b>Rate/PHI for foliar application.</b>
	<b>acetamiprid Assail 70WP</b>	<b>3.4 oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4A</b>	<b>1</b>	
	<b>buprofezin Centaur WDG</b>	<b>34.5 oz</b>	<b>12 h</b>	<b>14 d</b>	<b>16</b>	<b>NR</b>	
	<b>phosmet Imidan 70W</b>	<b>4-5 lb</b>	<b>7 d</b>	<b>7 d</b>	<b>1B</b>	<b>2</b>	
<b>Green apple aphid</b>	<b>acetamiprid Assail 70WP</b>	<b>1.7 oz</b>	<b>12 h</b>	<b>7 d</b>	<b>4A</b>	<b>3-4</b>	
<b>Lacanobia fruitworm</b>	<b>indoxacarb Avaunt</b>	<b>3-6 oz</b>	<b>12 h</b>	<b>14 d</b>	<b>22</b>	<b>4</b>	

<i>Insect</i>	<i>Chemical</i>	<i>Rate per Acre</i>	<i>REI</i>	<i>PHI</i>	<i>MOA</i>	<i>Eff.</i>	<i>Notes</i>
	<i>spinosad</i> <i>Entrust</i>	<i>3 oz</i>	<i>4 h</i>	<i>7 d</i>	<i>5</i>	<i>3-4</i>	<i>Entrust is a spinosad formulation registered for organic apple production. It is effective against codling moth larvae hatching from the egg. It has a residual activity of 7 to 10 days. When Entrust is incorporated into an organic pest control program using pheromones, summer oil, and codling moth virus, good control of this key pest is possible. Some leafroller populations have developed resistance to Success (spinosad) and its use could result in reduced levels of control. Success is in the same chemical class as Delegate (spinetoram) so avoid using these products against two consecutive generations.</i> <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<i>methoxyfenozide</i> <i>Intrepid 2F</i>	<i>16 fl oz</i>	<i>4 h</i>	<i>14 d</i>	<i>18A</i>	<i>3-4</i>	
	<i>spinosad</i> <i>Success</i>	<i>6-10 fl oz</i>	<i>4 h</i>	<i>7 d</i>	<i>5</i>	<i>3-4</i>	
	<i>kaolin clay</i> <i>Surround WP</i>	<i>50 lb</i>	<i>4 h</i>	<i>0 d</i>		<i>3-4</i>	<span style="border: 1px solid black; padding: 2px;">Organic</span>
<i>Leafrollers (Pandemis, Obliquebanded)</i>	<i>chlorantraniliprole</i> <i>Altacor</i>	<i>3-4.5 oz</i>	<i>4 h</i>	<i>5 d</i>	<i>28</i>	<i>4</i>	<i>Altacor is very effective against leafroller larvae and codling moth eggs and larvae. It has a residual activity of up to 17 days when using the high label rate or not more than 14 days when using the low label rate. Altacor is in the same chemical class as Belt.</i>
	<i>spinetoram</i> <i>Delegate WG</i>	<i>4.5-7 oz</i>	<i>4 h</i>	<i>7 d</i>	<i>5</i>	<i>4</i>	<i>Delegate is very effective against leafroller and codling moth larvae. It has a residual activity of 14 days. Delegate is in the same chemical class as Success (spinosad) so avoid using these products against two consecutive generations.</i>
	<i>Bacillus thuringiensis subsp. kurstaki</i> <i>DiPel DF</i>	<i>See label</i>	<i>4 h</i>	<i>0 d</i>	<i>11B2</i>	<i>3</i>	<i>Two or three Bt applications are usually required to achieve acceptable control. Time the first application to coincide with leafroller egg hatch. A repeat application might be required if leafroller populations are high.</i> <span style="border: 1px solid black; padding: 2px;">Organic</span>

<i>Insect</i>	<i>Chemical</i>	<i>Rate per Acre</i>	<i>REI</i>	<i>PHI</i>	<i>MOA</i>	<i>Eff.</i>	<i>Notes</i>
	<i>spinosad Entrust</i>	<i>3 oz</i>	<i>4 h</i>	<i>7 d</i>	<i>5</i>	<i>3-4</i>	<i>Entrust is a spinosad formulation registered for organic apple production. It is effective against codling moth larvae hatching from the egg. It has a residual activity of 7 to 10 days. When Entrust is incorporated into an organic pest control program using pheromones, summer oil, and codling moth virus, good control of this key pest is possible. Some leafroller populations have developed resistance to Success (spinosad) and its use could result in reduced levels of control. Success is in the same chemical class as Delegate (spinetoram) so avoid using these products against two consecutive generations.</i> <span style="border: 1px solid black; padding: 2px;">Organic</span>
	<i>cyantraniliprole Exirel</i>	<i>10-17 fl oz</i>	<i>12 h</i>	<i>3 d</i>	<i>28</i>	<i>4</i>	
	<i>methoxyfenozide Intrepid 2F</i>	<i>16 fl oz</i>	<i>4 h</i>	<i>14 d</i>	<i>18A</i>	<i>3</i>	
	<i>emamectin benzoate Proclaim</i>	<i>3.2-4.8 oz</i>	<i>12 h   48 h for some activities-see label</i>	<i>14 d</i>	<i>6</i>	<i>4</i>	
	<i>novaluron Rimon 0.83EC</i>	<i>30-50 fl oz</i>	<i>12 h</i>	<i>14 d</i>	<i>15</i>	<i>3-4</i>	
	<i>spinosad Success</i>	<i>6-10 fl oz</i>	<i>4 h</i>	<i>7 d</i>	<i>5</i>	<i>3-4</i>	
<i>San Jose scale</i>	<i>diazinon Diazinon 50W</i>	<i>4 lb</i>	<i>4 d</i>	<i>21 d</i>	<i>1B</i>	<i>3</i>	
<i>Shothole borer</i>	<i>esfenvalerate Asana XL</i>	<i>14.5 fl oz</i>	<i>12 h</i>	<i>21 d</i>	<i>3A</i>	<i>4</i>	
	<i>acetamiprid Assail 70WP</i>	<i>3.4 oz</i>	<i>12 h</i>	<i>7 d</i>	<i>4A</i>	<i>NR</i>	
<i>Spider mites</i>	<i>bifenazate Acramite 50WS</i>	<i>0.75-1 lb</i>	<i>12 h</i>	<i>7 d</i>	<i>un</i>	<i>4</i>	
	<i>spirodiclofen Envidor 2SC</i>	<i>16-18 fl oz</i>	<i>12 h</i>	<i>7 d</i>	<i>23</i>	<i>3-4</i>	
	<i>fenpyroximate FujiMite XLO</i>	<i>1-2 pt</i>	<i>12 h</i>	<i>14 d</i>	<i>21A</i>	<i>3-4</i>	

<b>Insect</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
	<i>pyridaben</i> Nexter 75WSB	4.4-8.8 oz	12 h	25 d	21A	2-3	Use a low to moderate rate for European red mite; a moderate to high rate for twospotted and McDaniel spider mites.
	<i>hexythiazox</i> Savey 50DF	4-6 oz	12 h	28 d	10A	NR	Most effective on the egg stage. When mite populations are high and leaf bronzing has already occurred, a miticide effective on the adult stage may be used in combination.
	<i>etoxazole</i> Zeal Miticide1 72WSP	2-3 oz	12 h	14 d	10B	3-4	
<b>Western tentiform leafminer</b>	<i>abamectin</i> Agri-Mek SC	4.25 fl oz	12 h	28 d	6	4	May provide control of mites if used early in the season.
	<i>spinosad</i> Success	6-10 fl oz	4 h	7 d	5	3-4	
<b>White apple leafhopper</b>	<i>imidacloprid</i> Admire Pro	1.4-2.8 fl oz	12 h	7 d	4A	4	Do not use until pollination is complete and bees have been removed from the area. Rate indicated is for foliar application.
	<i>indoxacarb</i> Avaunt	4-6 oz	12 h	14 d	22	3-4	
	<i>carbaryl</i> Sevin XLR Plus	1-2 qt	12 h	3 d	1A	NR	Carbaryl may disrupt integrated mite control depending on history of use. Use higher rate if leafhopper population is comprised primarily of adults.
	<i>kaolin clay</i> Surround WP	See label	4 h	0 d		NR	Organic
	<b>Notes: Carbaryl may disrupt integrated mite control depending on history of use. Use higher rate if leafhopper population is composed primarily of adults.</b>						
<b>Woolly apple aphid</b>	<i>sulfoxaflor</i> Closer SC	4.25-5.75 fl oz	12 h	7 d	4C	3	Use an adjuvant such as Exit at 0.25% vol:vol.
	<i>diazinon</i> Diazinon 50W	4 lb	4 d	21 d	1B	4	
	<i>spirotetramat</i> Ultror	10-14 fl oz	24 h	7 d	23	2-4	Ultror may only provide suppression if used in mid- and late-season sprays; preliminary evidence suggests that better efficacy may be obtained when it is applied prophylactically shortly after petal fall.
<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
Apple mildew	flutianil Gatten Fungicide	8 fl oz	12 h	14 d	U13	NR	The mechanism of action of Gatten is yet unknown. Rotate with fungicides from different chemical groups to reduce fungicide resistance.
	potassium bicarbonate Kaligreen	3 lb	4 h	1 d	NC	2	Under low disease pressure, Kaligreen will provide adequate control. If disease pressure increases, rotation or combination with other materials is recommended. <span>Organic</span>
	fluopyram Luna Privilege	6.84 fl oz	12 h	7 d	7	4	Luna Privilege is a FRAC group 7 fungicide.
	fluxapyroxad + pyraclostrobin Merivon	5.5 fl oz	12 h	0 d	7, 11	3	Merivon is a FRAC group 7 fungicide.
	polyoxin D zinc salt OSO 5%SC	13 fl oz	4 h	0 d	19	3	Do not apply more than 4.2 oz. a.i./acre/season.
	polyoxin D zinc salt Ph-D	6.2 oz	4 h	0 d	19	NR	Ph-D is similar to OSO5%SC. Use only one of these fungicides in a given season.
	pyraclostrobin + boscalid Pristine	18.5 oz	12 h	0 d	11,7	3	Pristine is a FRAC 7 +11 fungicide and should not be rotated or used with other FRAC 7+11 fungicides. The efficacy level will depend on the occurrence of fungicide resistant populations of the pathogen in the orchards. For powdery mildew, preferably use other FRAC 7 fungicides in spring.
	Reynoutria sachalinensis Regalia	4 qt	4 h	0 d	P5	2	Do not use prior to petal fall. Under low disease pressure, it may help control some summer diseases like Alternaria and Bull's eye rot. <span>Organic</span>
	triflumazole Trionic 4SC	16 fl oz	12 h	14 d	3	4	Trionic is a FRAC 3 fungicide and should not be rotated or used with other FRAC 3 fungicides.
Apple scab	captan Captan 50WP	8 lb	24 h	0 d	M4	NR	
	trifloxystrobin Flint Extra	2.5-2.9 fl oz	12 h	14 d	11	3	
	fluopyram + trifloxystrobin Luna Sensation	4-5.8 fl oz	12 h	14 d	7, 11	NR	Luna Sensation and Fontelis are from the same Chemical group (7). Use only one of them.
	triflumizole Procure 480SC	8-16 fl oz	12 h	14 d	3	NR	Rates vary with postinfective schedule; see label. Apply only one Group 3 Fungicide at the same growth stage.
	myclobutanil Rally 40WSP	5 oz	24 h	14 d	3	NR	Use only one Group 3 fungicide at the same growth stage.

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
	<i>flutriafol</i> Topguard	13 fl oz	12 h	14 d	3	NR	Topguard may cause a mild phytotoxicity on Braeburn apple leaves.
Notes: To limit the potential for development of fungicide resistance, do not make more than four applications of strobilurin fungicides per season. Do not make more than two sequential applications of strobilurin fungicides. This limitation is inclusive of all strobilurin fungicides (Mode 11) labeled for use on pome fruits.							
Bull's eye rot	<i>ziram</i> Ziram Granuflo 76WDG	6 lb	48 h	14 d	M3	3	Apply 12 to 14 days after calyx stage and repeat as needed through the summer. Do not apply more than 24.2 lbs per acre in a crop cycle. Do not apply within 14 days of harvest. Tank-mixing with other fungicides has been reported to increase efficacy and reduce fungicide resistance development.
Pathogens of Storage Rots	<i>Reynoutria sachalinensis</i> Regalia	4 qt	4 h	0 d	P5	2	Do not use prior to petal fall. Under low disease pressure, it may help control some summer diseases like <i>Alternaria</i> and Bull's eye rot. <span style="border: 1px solid black; padding: 2px;">Organic</span>

Efficacy numbers denote the relative efficacy of a pesticide against a given pest on a 1 to 4 scale with 1 being low and 4 high efficacy. This information is based primarily on research conducted with WSU researchers in Washington.

## Preharvest

Disease	Chemical	Rate per Acre	REI	PHI	MOA	Eff.	Notes
Apple scab	<i>captan</i> Captan 50WP	6 lb	24 h	0 d	M4	NR	When possible, tank-mix captan with other single-site fungicides to help reduce fungicide resistance development.
Botrytis-Gray Mold	<i>copper octanoate</i> Cueva	8 qt	4 h	0 d	M1	2	<span style="border: 1px solid black; padding: 2px;">Organic</span>
	<i>thiophanate-methyl</i> Topsin M WSB	1 lb	2 d	1 d	1	3	Efficacy level is ensured when resistance is absent from the orchard. Do not apply more than 2.8 lbs. a.i. per acre in a year. Topsin-M is very prone to fungicide resistance development, rotation with other FRAC group is required. Do not make sequential applications of FRAC group 1 fungicides and do not make more than two FRAC 1 fungicides per season. Topsin-M is similar to Mertect, used postharvest, therefore careful use is highly recommended.

<b>Disease</b>	<b>Chemical</b>	<b>Rate per Acre</b>	<b>REI</b>	<b>PHI</b>	<b>MOA</b>	<b>Eff.</b>	<b>Notes</b>
<b>Bull's eye rot</b>	<b>captan Captan 50WP</b>	<b>6 lb</b>	<b>24 h</b>	<b>0 d</b>	<b>M4</b>	<b>NR</b>	<b>Do not apply more than 64 lbs. of Captan 50 Wettable Powder per acre per crop cycle. Make 1 or 2 applications with late cover sprays and 1 final spray prior to harvest. When possible, tank-mix captan with other single-site fungicides to help reduce fungicide resistance development.</b>
	<b>thiophanate-methyl Topsin M WSB</b>	<b>1 lb</b>	<b>2 d</b>	<b>1 d</b>	<b>1</b>	<b>4</b>	<b>Efficacy level is ensured when resistance is absent from the orchard. Do not apply more than 2.8 lbs. a.i. per acre in a year. Topsin-M is very prone to fungicide resistance development, rotation with other FRAC group is required. Do not make sequential applications of FRAC group 1 fungicides and do not make more than two FRAC 1 fungicides per season. Topsin-M is similar to Mertect, used postharvest, therefore careful use is highly recommended.</b>
	<b>ziram Ziram 76DF</b>	<b>See label</b>	<b>48 h</b>	<b>14 d</b>	<b>M3</b>	<b>3</b>	<b>Apply in the first cover spray and in preharvest ideally before rain. Do NOT apply within 14 days of harvest. When applicable, tank-mix with other single-site fungicides to increase efficacy and reduce risk of fungicide resistance development. Do not apply more than 18.4 lbs. a.i of Ziram 76DF or equivalent per season.</b>
<b>Pathogens of Storage Rots</b>	<b>hydrogen peroxide + peroxyacetic acid Jet-Ag</b>	<b>0.8 - 3.9 fl oz</b>	<b>4 h</b>	<b>none listed</b>		<b>3</b>	
	<b>fluxapyroxad + pyraclostrobin Merivon</b>	<b>5.5 fl oz</b>	<b>12 h</b>	<b>0 d</b>	<b>7, 11</b>	<b>3</b>	<b>Merivon is a FRAC group 7 fungicide and should not be rotated or used with other FRAC 7 fungicides. Do not make more than three FRAC 7 applications in a season.</b>
	<b>polyoxin D zinc salt OSO 5%SC</b>	<b>13 fl oz</b>	<b>4 h</b>	<b>0 d</b>	<b>19</b>	<b>3</b>	<b>OSO will help control Alternaria infections preharvest. OSO is FRAC 19 fungicide recommended to rotate with other FRAC groups labeled for pome fruit. Do not apply more than 4.2 oz. a.i./acre/season.</b>
	<b>polyoxin D zinc salt Ph-D</b>	<b>6.2 oz</b>	<b>4 h</b>	<b>0 d</b>	<b>19</b>	<b>NR</b>	<b>Ph-D is similar to OSO 5%, so use only one of them in a given season.</b>



<i>Disease</i>	<i>Chemical</i>	<i>Rate per Acre</i>	<i>REI</i>	<i>PHI</i>	<i>MOA</i>	<i>Eff.</i>	<i>Notes</i>
	<i>pyraclostrobin + boscalid Pristine</i>	14.5-18.5 oz	12 h	0 d	11,7	3	<i>Pristine is a FRAC 7 +11 fungicide an should not be rotated or used with other FRAC 7+11 fungicides. To limit the potential for development of fungicide resistance, do not make more than 3 applications of FRAC 7+11 fungicides per season. Do not make sequential applications of FRAC 7+11 fungicides labeled for use on pome fruits. The efficacy level will depends on the occurrence of fungicide resistant populations of the pathogen in the orchards. For powdery mildew, preferably use other FRAC 7 fungicides in spring.</i>
	<i>thiophanate-methyl Topsin M WSB</i>	1 lb	2 d	1 d	1	3	<i>Efficacy level is ensured when resistance is absent from the orchard. Do not apply more than 2.8 lbs. a.i. per acre in a year. Topsin-M is very prone to fungicide resistance development, rotation with other FRAC group is required. Do not make sequential applications of FRAC group 1 fungicides and do not make more than two FRAC 1 fungicides per season. Topsin-M is similar to Mertect, used postharvest, therefore careful use is highly recommended.</i>
	<i>ziram Ziram Granuflo 76WDG</i>	6 lb	48 h	14 d	M3	2	<i>When applicable, tank-mix with other single-site fungicides to reduce risk fo fungicide resistance development.</i>
<i>Insect</i>	<i>Chemical</i>	<i>Rate per Acre</i>	<i>REI</i>	<i>PHI</i>	<i>MOA</i>	<i>Eff.</i>	<i>Notes</i>
<i>Apple maggot</i>	<i>thiamethoxam Actara</i>	5.5 oz	12 h	14 d/35 d	4A	NR	<i>Estimated residual activity: 7-14 days.</i>
	<i>chlorantraniliprole Altacor</i>	4.5 oz	4 h	5 d	28	NR	<i>Provides suppression only. Also effective against codling moth and leafrollers. Estimated residual activity: 10-14 days.</i>
	<i>acetamiprid Assail 70WP</i>	3.4 oz	12 h	7 d	4A	NR	<i>Assail is also used to control codling moth. The estimated residual activity is 10-14 days.</i>
	<i>fenpropathrin Danitol 2.4EC</i>	16-21.3 fl oz	24 h	14 d	3	NR	<i>Also effective against codling moth and leafrollers. Estimated residual activity: 7-10 days.</i>
	<i>spinetoram Delegate WG</i>	6-7 oz	4 h	7 d	5	NR	<i>Provides suppression only. Also effective against codling moth and leafrollers. Estimated residual activity: 10-14 days.</i>

<i>Insect</i>	<i>Chemical</i>	<i>Rate per Acre</i>	<i>REI</i>	<i>PHI</i>	<i>MOA</i>	<i>Eff.</i>	<i>Notes</i>
	<i>spinosad Entrust</i>	<i>2-3 oz</i>	<i>4 h</i>	<i>7 d</i>	<i>5</i>	<i>NR</i>	<i>Provides suppression only. Also effective against codling moth and leafrollers. Estimated residual activity: 5-7 days. <span>Organic</span></i>
	<i>phosmet Imidan 70W</i>	<i>5.33 lb</i>	<i>7 d</i>	<i>7 d</i>	<i>1B</i>	<i>NR</i>	<i>Imidan is also used to control codling moth. The estimated residual activity is 14 days.</i>
	<i>lambda-cyhalothrin Warrior II</i>	<i>1.28-2.56 fl oz</i>	<i>24 h</i>	<i>21 d</i>	<i>3</i>	<i>NR</i>	<i>Also effective against codling moth and leafrollers. Estimated residual activity: 7-10 days.</i>
<i>Codling moth</i>	<i>acetamiprid Assail 70WP</i>	<i>3.4 oz</i>	<i>12 h</i>	<i>7 d</i>	<i>4A</i>	<i>NR</i>	
	<i>spinetoram Delegate WG</i>	<i>6-7 oz</i>	<i>4 h</i>	<i>7 d</i>	<i>5</i>	<i>4</i>	<i>Delegate is effective against codling moth larvae. It has a residual activity of 14 days.</i>
	<i>spinosad Entrust SC</i>	<i>3 fl oz</i>	<i>4 h</i>	<i>7 d</i>	<i>5</i>	<i>NR</i>	<i>Entrust is a spinosad formulation registered for organic apple production. It is effective against codling moth larvae hatching from the egg. It has a residual activity of 7 to 10 days. When Entrust is incorporated into an organic pest control program using pheromones, summer oil, and codling moth virus, good control of this key pest is possible. Best results occur when applications are timed for egg hatch, which may occur during bloom. <span>Organic</span></i>
	<i>carbaryl Sevin XLR Plus</i>	<i>1-3 qt</i>	<i>12 h</i>	<i>3 d</i>	<i>1A</i>	<i>2</i>	
	<i>petroleum oil, summer petroleum oil, summer</i>	<i>See label</i>	<i>4 h</i>	<i>0 d</i>		<i>NR</i>	<span>Organic</span>
<i>Leafrollers (Pandemis, Obliquebanded)</i>	<i>Bacillus thuringiensis subsp. kurstaki DiPel DF</i>	<i>See label</i>	<i>4 h</i>	<i>0 d</i>	<i>11B2</i>	<i>3</i>	<i>Bts are stomach poisons so complete coverage is very important for control. Two or three applications are usually required. <span>Organic</span></i>

Efficacy numbers denote the relative efficacy of a pesticide against a given pest on a 1 to 4 scale with 1 being low and 4 high efficacy. This information is based primarily on research conducted with WSU researchers in Washington.